

#### Chapter 2-Part III (Sec 2.4)

#### **Basic Structures:**

- Sets
- Functions
- Sequences and Sums
- Cardinality of Sets

# Section 2.4

# **Section Summary**

- Sequences.
  - Examples: Geometric Progression, Arithmetic Progression
- Recurrence Relations
  - Example: Fibonacci Sequence
- Summations

#### Introduction

- Sequences are ordered lists of elements.
  - 1, 2, 3, 5, 8
  - 1, 3, 9, 27, 81, .....
- Sequences arise throughout mathematics, computer science, and in many other disciplines, ranging from botany to music.
- We will introduce the terminology to represent sequences and sums of the terms in the sequences.

#### Sequences

- **Definition**: A *sequence* is a function *f* from a subset of the integers to a set *S*.
  - Domain is usually either the set {0, 1, 2, 3, 4, ....} or {1, 2, 3, 4, ....}
  - *a<sub>n</sub>* is a *term* of the sequence, denotes the image of the integer *n*

• 
$$f(n) = a_n$$

#### Sequences

#### **Example**: Consider the sequence $\{a_n\}$ where

$$a_n = \frac{1}{n}$$
  $\{a_n\} = \{a_1, a_2, a_3, \ldots\}$ 

$$1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4} \dots$$

## **Geometric Progression**

**Definition**: A *geometric progression* is a sequence of the form:  $a, ar, ar^2, \ldots, ar^n, \ldots$ 

where the *initial term a* and the *common ratio r* are real numbers.

#### **Examples**:

1. Let a = 1 and r = -1. Then:

$$\{b_n\} = \{b_0, b_1, b_2, b_3, b_4, \dots\} = \{1, -1, 1, -1, 1, \dots\}$$

2. Let a = 2 and r = 5. Then:

$$\{c_n\} = \{c_0, c_1, c_2, c_3, c_4, \dots\} = \{2, 10, 50, 250, 1250, \dots\}$$

3. Let a = 6 and r = 1/3. Then:  $\{d_n\} = \{d_0, d_1, d_2, d_3, d_4, \dots\} = \{6, 2, \frac{2}{3}, \frac{2}{9}, \frac{2}{27}, \dots\}$ 

# **Arithmetic Progression**

**Definition**: A *arithmetic progression* is a sequence of the form:  $a, a + d, a + 2d, \dots, a + nd, \dots$ 

where the *initial term a* and the *common difference d* are real numbers.

**Examples**:

1. Let a = -1 and d = 4:

$$\{s_n\} = \{s_0, s_1, s_2, s_3, s_4, \dots\} = \{-1, 3, 7, 11, 15, \dots\}$$

2. Let a = 7 and d = -3:

$$\{t_n\} = \{t_0, t_1, t_2, t_3, t_4, \dots\} = \{7, 4, 1, -2, -5, \dots\}$$

3. Let a = 1 and d = 2:  $\{u_n\} = \{u_0, u_1, u_2, u_3, u_4, \dots\} = \{1, 3, 5, 7, 9, \dots\}$ 

# Strings

- **Definition**: A *string* is a finite sequence of characters from a finite set (an alphabet).
- Sequences of characters or bits are important in computer science.
- The *empty string* is represented by  $\lambda$ .
- The string *abcde* has *length* 5.

### **Recurrence Relations**

- **Definition:** A *recurrence relation* for the sequence  $\{a_n\}$  is an equation *that expresses*  $a_n$  *in terms of* one or more of the *previous terms of the sequence*, namely,  $a_0, a_1, ..., a_{n-1}$ , for all integers n with  $n \ge n_0$ , where  $n_0$  is a nonnegative integer.
- A sequence is called a *solution* of a recurrence relation if its terms satisfy the recurrence relation.
- The *initial conditions* for a sequence specify the terms that precede the first term where the recurrence relation takes effect.

#### Questions

**Example** 1: Let  $\{a_n\}$  be a sequence that satisfies the recurrence relation  $a_n = a_{n-1} + 3$  for n = 1,2,3,4,... and suppose that  $a_0 = 2$ . What are  $a_1$ ,  $a_2$  and  $a_3$ ?

-Note that  $a_0 = 2$  is the initial condition.

**Solution**: From the recurrence relation :

$$a_1 = a_0 + 3 = 2 + 3 = 5$$
  
 $a_2 = 5 + 3 = 8$   
 $a_3 = 8 + 3 = 11$ 

#### Questions

**Example** 2: Let  $\{a_n\}$  be a sequence that satisfies the recurrence relation  $a_n = a_{n-1} - a_{n-2}$  for n = 2,3,4,... and suppose that  $a_0 = 3$  and  $a_1 = 5$ . What are  $a_2$  and  $a_3$ ?

**Solution**: From the recurrence relation:

$$a_2 = a_1 - a_0 = 5 - 3 = 2$$
  
 $a_3 = a_2 - a_1 = 2 - 5 = -3$ 

## Fibonacci Sequence

**Definition**: Define the *Fibonacci sequence*,  $f_0$ ,  $f_1$ ,  $f_2$ , ..., by:

- Initial Conditions:  $f_0 = 0, f_1 = 1$
- Recurrence Relation:  $f_n = f_{n-1} + f_{n-2}$

**Example**: Find  $f_2, f_3, f_4, f_5$  and  $f_6$ .

#### **Answer:**

$$\begin{split} f_2 &= f_1 + f_0 = 1 + 0 = 1, \\ f_3 &= f_2 + f_1 = 1 + 1 = 2, \\ f_4 &= f_3 + f_2 = 2 + 1 = 3, \\ f_5 &= f_4 + f_3 = 3 + 2 = 5, \\ f_6 &= f_5 + f_4 = 5 + 3 = 8. \end{split}$$

# **Solving Recurrence Relations**

- Finding a formula for the *n*th term of the sequence generated by a recurrence relation is called *solving the recurrence relation*.
- Such a formula is called a *closed formula*.
- Various methods for solving recurrence relations will be covered in Chapter 8 where <u>recurrence relations will be</u> <u>studied in greater depth</u>.
- Here we illustrate by example the method of iteration in which we need to guess the formula. The guess can be proved correct by the method of induction (Chapter 5).

### **Iterative Solution Example**

**Method** 1: Working upward, forward substitution Let  $\{a_n\}$  be a sequence that satisfies the recurrence relation  $a_n = a_{n-1} + 3$  for n = 2,3,4,... and suppose that  $a_1 = 2$ .

$$a_{2} = 2 + 3$$

$$a_{3} = (2 + 3) + 3 = 2 + 3 \cdot 2$$

$$a_{4} = (2 + 2 \cdot 3) + 3 = 2 + 3 \cdot 3$$

$$\vdots$$

$$a_{n} = a_{n-1} + 3 = (2 + 3 \cdot (n - 2)) + 3 = 2 + 3(n - 1)$$

#### **Iterative Solution Example**

**Method** 2: Working downward, backward substitution Let  $\{a_n\}$  be a sequence that satisfies the recurrence relation  $a_n = a_{n-1} + 3$  for n = 2,3,4,... and suppose that  $a_1 = 2$ .