

BBM205 Midterm Exam II

Time: 15:45-17:30

30 December 2020

Question:	1	2	3	4	5	6	7	Total
Points:	15	22	12	16	10	10	15	100
Score:								

1. (15 points) Determine whether the following statements are true or false.

Solution: If a graph has a triangle as a subgraph, then its vertex set cannot be properly colored using 2 colors only.

(X) True () False

If a graph on n vertices has a Hamilton cycle, then the minimum degree of this graph is at least $n/2$.

() True (X) False

If G has no subgraph homeomorphic to K_5 or $K_{3,3}$, then it is planar.

(X) True () False

The average degree of every planar graph is less than 6.

(X) True () False

A (multi)graph with degree list 2,2,4,4,4 has an Euler circuit.

(X) True () False

2. Suppose that you have n cards that are numbered 1 through n in a box. You pick a card uniformly at random, then put it back into the box. And you pick a second card uniformly at random, and put it back into the box. Finally, you pick a third card uniformly at random.

You can leave your answers for the following questions as an expression of n . Show your work to receive full credit.

(a) (4 points) What is the probability that all three numbers are different?

(b) (5 points) What is the probability that a number repeated exactly twice?

(c) (6 points) What is the probability that the three numbers on the cards are strictly increasing?

- (d) (7 points) What is the probability that the sum of the numbers on the cards add up to n ?

Solution:

- (a) (4 points)

$$\frac{\binom{n}{3}3!}{n^3} = \frac{n(n-1)(n-2)}{n^3}$$

- (b) (5 points)

$$\frac{\binom{n}{2} \cdot 2 \cdot \binom{3}{2}}{n^3} = \frac{3n(n-1)}{n^3}$$

- (c) (6 points)

$$\frac{\binom{n}{3} \cdot 1}{n^3} = \frac{n(n-1)(n-2)}{6n^3}$$

- (d) (7 points) The number of possibilities for the numbers on the cards is given by the set of solutions for $x_1 + x_2 + x_3 = n$ with $x_i \geq 1$.

$$\frac{\binom{(n-3)+2}{2}}{n^3} = \frac{(n-1)(n-2)}{2n^3}$$

3. (12 points) Let n be a nonnegative integer. By considering the number of solutions with nonnegative terms for

$$x_1 + x_2 + \cdots + x_k = n$$

and for

$$x_1 + x_2 + \cdots + x_k \leq n,$$

prove the following identity using a **combinatorial proof**.

$$\sum_{i=0}^n \binom{i+k-1}{k-1} = \binom{n+k}{k}$$

Solution: In the sum, the number of nonnegative solutions for $x_1 + x_2 + \dots + x_k = i$ are added. For each solution, we can define an auxiliary (slack) variable as $y = n - i$. Hence, the formula on the left gives the number of nonnegative solutions for $x_1 + x_2 + \dots + x_k + y = n$. This also equals the expression on the right.

4. Let a be your group number. For a coin, we have $\Pr(H)=a/5$ and $\Pr(T)=(5-a)/5$. Let X be the number of H's in a sequence of 10 flips using this coin. Answer the following with a formula without evaluating the answer.

- (a) (5 points) $Pr(\text{number of H's} > \text{number of T's})$
 (b) (5 points) $Pr(X \geq 3)$
 (c) (6 points) $Pr(X = 3 | \text{First flip is H})$

Solution:

- (a) (5 points) We add up the probabilities for the events, where $X = 6, 7, 8, 9, 10$.

$$\sum_{i=6}^{10} \binom{10}{i} \left(\frac{a}{5}\right)^i \left(\frac{5-a}{5}\right)^{10-i}$$

- (b) (5 points)

$$\sum_{i=3}^{10} \binom{10}{i} \left(\frac{a}{5}\right)^i \left(\frac{5-a}{5}\right)^{10-i}$$

- (c) (6 points) Let A be the event $X = 3$ and B the event where the first flip is H.

$$Pr(A|B) = \frac{Pr(A \cap B)}{Pr(B)} = \frac{\binom{9}{2} (a/5)^3 ((5-a)/5)^7}{a/5}.$$

5. (10 points) A bag contains 8 green and 6 purple socks, all unmatched. A man picks the socks without looking at them.

Solution:

- (a) (5 points) Mark the minimum number of socks that he must pick to guarantee having two socks of the same color:

2 3 4 5

- (b) (5 points) Mark the minimum number of socks that he must pick to guarantee having two *purple* socks:

3 6 8 10

6. (a) (5 points) **Solution:** What is the coefficient of $x^{11}y^8$ in $(3x - 5y)^{19}$?

$$C(19, 11)C(19, 8)3^{11}(-5)^8$$

$$C(19, 8)3^8(-5)^{11}$$

(X)	$C(19, 11)3^{11}(-5)^8$
()	$C(19, 11)(-15)^{19}$

(b) (5 points) **Solution:** There are 12 physics and 9 literature books on a shelf. How many ways are there to order them so that no two literature books are next to each other?

()	$P(21, 12) \cdot 9!$
(X)	$P(13, 9) \cdot 12!$
()	$C(20, 8) \cdot 12!9!$
()	$P(21, 9) \cdot 12!$

7. Show your work to receive full credit.

Let b_n be the number of bijections $f : A \rightarrow A$, where $A = \{1, 2, \dots, n\}$ and $f(i) \neq i$ for all i .

(a) (5 points) Write the number of bijections f , for which $f(1) = k$ and $f(k) = 1$ for some $k \neq 1$, in terms of b_{n-1}, b_{n-2}, \dots

Solution: There are b_{n-2} of them, considering that we are looking for bijections $g : B \rightarrow B$, where $B = A - \{1, k\}$ and $f(i) \neq i$ for all $i \in B$. Since there are $n - 1$ possible values for k , the total number is $(n - 1)b_{n-2}$.

(b) (5 points) Write the number of bijections f , for which $f(1) = k$ and $f(k) \neq 1$ for some $k \neq 1$, in terms of b_{n-1}, b_{n-2}, \dots

Solution: We consider the bijections $h : C \rightarrow C$, where $C = A - \{k\}$, with $h(i) \neq i$ and then we let $f(k) = h(1)$ to obtain a bijection $f : A \rightarrow A$ with $f(1) = k$. Hence, there are b_{n-1} of these. Since there are $n - 1$ possible values for k , the total number is $(n - 1)b_{n-1}$.

(c) (5 points) Use the previous two parts to obtain a recursive formula for b_n .

Solution: The set of bijections in parts (a) and (b) do not intersect. Hence, if we add up these numbers and consider $n - 1$ different values for k , we have all bijections counted by b_n . So, $b_n = (n - 1)(b_{n-1} + b_{n-2})$.