

BBM205

Fall 2022-2023, Final Exam

January 18, 2023, 15:00-16:40

Question:	1	2	3	4	5	6	7	8	Total
Points:	13	14	12	13	10	15	15	8	100
Score:									

Name, Surname, ID: _____ Instructor: _____

1. (13 points) Let x be a prime number and $y > 1$ be an integer. **Prove by contradiction** that $x \cdot y$ is not a prime number.

Solution: Let $P(x, y)$ be the predicate "x is a prime number and y is an integer that is greater than 1" and let $Q(x, y)$ be the predicate "xy is not a prime number." We will try to prove $P(x, y) \rightarrow Q(x, y)$ using contradiction.

To prove using contradiction, we will assume that $P(x, y)$ is true and $\neg Q(x, y)$ is also true. In other words, we assume that x is a prime number and $y > 1$ and xy is a prime number. xy can be prime only if $y = 1$ (since x is prime), which contradicts with the premise of this implication ($P(x, y)$), which states that $y > 1$. Since our assumption leads to a contradiction ($P(x, y)$ is false), $P(x, y)$ and $\neg Q(x, y)$ can not both be true. So, we conclude that $P(x, y) \rightarrow Q(x, y)$ must be true.

2. (a) (7 points) How many ordered pairs of integers (a, b) are needed (at minimum) to guarantee two pairs (a_1, b_1) and (a_2, b_2) such that $a_1 \equiv a_2 \pmod{7}$ and $b_1 \equiv b_2 \pmod{7}$?

Solution: There are $7^2 = 49$ distinct pairs as $A \times A$, where $A = \{0, 1, \dots, 6\}$. By pigeonhole principle, having 50 pairs guarantees to have one pair repeat at least once. Here, we assume that the pairs are the pigeons and 49 different possibilities are the pigeonholes.

- (b) (7 points) Show that if $f : S \rightarrow T$ is a function, where S and T are finite sets with $|S| > |T|$, then f is not injective (one-to-one).

Solution: Let $S = \{x_1, \dots, x_m\}$. Let the elements of $T = \{y_1, \dots, y_n\}$ be the pigeonholes, where $n > m$ and the values of the function $\{f(x_1), \dots, f(x_m)\}$ be the pigeons. Since, $n > m$, some y_i will be the value of f on at least two elements in S .

3. How many ways are there to put 12 balls into 9 boxes, assuming:

- (a) (4 points) the balls and the boxes are distinguishable (different)?

Solution: 9^{12}

- (b) (4 points) the balls are identical and the boxes are distinguishable?

Solution: $\binom{12+9-1}{9-1} = \binom{20}{8}$

- (c) (4 points) the balls and the boxes are identical, and none of the boxes are left empty?

Solution: There are three different ways to distribute the balls into 9 boxes: $\{4, 1, \dots, 1\}$, $\{3, 2, 1, \dots, 1\}$, $\{2, 2, 2, 1, \dots, 1\}$.

4. Suppose you record the **months** of the birthdays of a large group of people, one at a time until you have found a **match**, that is, a **month** of a birthday that has already been recorded.

- (a) (4 points) What is the probability that after the first 5 people's birthday months are recorded, no match has occurred (that is, each person has a birthday in a different month)?

$$\text{Solution: } \frac{P(12,5)}{12^5}$$

- (b) (4 points) What is the probability that it takes more than 10 people for a match to occur?

$$\text{Solution: } \frac{P(12,10)}{12^{10}}$$

- (c) (5 points) Suppose instead that you record the birthday months of a large group of people, one at a time, until you have found a person whose birthday month matches your own birthday month. What is the probability that it takes exactly 10 people for this to occur?

$$\text{Solution: } \frac{11^9}{12^{10}}$$

5. Suppose there are 25 balls in a bag, consisting of 12 red, 8 blue and 5 green balls;

- (a) (5 points) What is the minimum number of balls a person must pick up from the bag (without replacing any of it) to guarantee that he has at least one ball of each color?

$$\text{Solution: } 12 + 8 + 1 = 21$$

- (b) (5 points) How many balls a person must pick to make sure he has at least two balls with the same color?

$$\text{Solution: } \text{Using generalized pigeonhole principle: } \lceil N/3 \rceil \geq 2 \\ N = 4$$

6. (15 points) In a certain city, the probability of a person having a disease is 0.1. The city's hospital has two tests for the disease, Test A and Test B. Test A is 95% accurate if the person has the disease and 90% accurate if the person does not have the disease. Test B is 92% accurate if the person has the disease and 85% accurate if the person does not have the disease. A person is selected at random from the city and tests positive for both Test A and Test B. What is the probability that the person actually has the disease? (Assume that Test A and Test B generate results independently.)

Solution: Using the Bayes' theorem:

$$P(\text{disease}|\text{positiveA}, \text{positiveB}) = \frac{P(\text{positiveA}, \text{positiveB}|\text{disease}) * P(\text{disease})}{P(\text{positiveA}, \text{positiveB})}$$

We know that $P(\text{disease}) = 0.1$ and

$$P(\text{positiveA}|\text{disease}) = 0.95, P(\text{negativeA}|\text{disease}) = 0.05$$

$$P(\text{positiveB}|\text{disease}) = 0.92, P(\text{negativeB}|\text{disease}) = 0.08$$

$$P(\text{positiveA}|\text{nodisease}) = 0.1, P(\text{negativeA}|\text{nodisease}) = 0.9$$

$$P(\text{positiveB}|\text{nodisease}) = 0.15, P(\text{negativeB}|\text{nodisease}) = 0.85$$

We can use these values to find $P(\text{positiveA}, \text{positiveB})$

$$\begin{aligned} P(\text{positiveA}, \text{positiveB}) &= P(\text{positiveA}|\text{disease}) * P(\text{positiveB}|\text{disease}) * P(\text{disease}) + \\ &P(\text{positiveA}|\text{nodisease}) * P(\text{positiveB}|\text{nodisease}) * P(\text{nodisease}) \\ &= 0.95 * 0.92 * 0.1 + 0.10 * 0.15 * 0.9 \\ &= 0.0874 + 0.0135 \\ &= 0.1009 \text{ (assume 0.1)} \end{aligned}$$

Substituting the values in Bayes' theorem:

$$P(\text{disease}|\text{positiveA}, \text{positiveB}) = \frac{P(\text{positiveA}|\text{disease}) * P(\text{positiveB}|\text{disease}) * P(\text{disease})}{P(\text{positiveA}, \text{positiveB})}$$

$$= 0.95 * 0.92 * 0.1 / 0.1$$

$$= 0.874$$

This will give the probability that a person who tests positive for both Test A and Test B actually has the disease.

7. (15 points) **Prove or disprove** the following claim: “For any three consecutive primes x, y, z given in ascending order (so, x is smallest and z is largest), $x + z \geq 2y$ ”.

Solution: The claim says $\forall(x, y, z) (x + z \geq 2y)$. To disprove the claim, we must show that the negation is true. So, we want to show that

$$\neg \forall(x, y, z) (x + z \geq 2y)$$

$$\exists(x, y, z) \neg(x + z \geq 2y)$$

$$\exists(x, y, z) (x + z < 2y)$$

Now, we can use some arithmagic:

$$\exists(x, y, z) (z - y < y - x)$$

So, we want to show that there exists three consecutive primes for which the difference between the smaller ones is greater than the difference between the larger ones. If we can find one example, we can use the Rule of Existential Generalization to show that $\exists(x, y, z) (z - y < y - x)$.

Consider (2, 3, 5): $(5 - 3 \not< 3 - 2)$ didn't work!

Consider (3, 5, 7): $(7 - 5 \not< 5 - 3)$ didn't work!

Consider (5, 7, 11): $(11 - 7 \not< 7 - 5)$ didn't work!

Consider (7, 11, 13): $(13 - 11 < 11 - 7)$ DID WORK!!!

So, for $x = 7, y = 11$ and $z = 13$ we show $(z - y < y - x)$ and therefore by the Rule of Existential Generalization

$\exists(x, y, z) (z - y < y - x)$. Hence, the claim is disproven.

8. Find the coefficient of the following terms:

(a) (4 points) x^7y^{12} in $(x + y)^{19}$

Solution: x^7y^{12} in $(x + y)^{19} = \binom{19}{7} = \binom{19}{12}$.

(b) (4 points) x^9y^4 in $(-3x - 5y)^{13}$

Solution: x^9y^4 in $(-3x - 5y)^{13} = \binom{13}{9}(-3)^9(-5)^4$

Name: