

# BBM205

Fall 2023-2024, Final Exam

January 10, 2024, 13:00-14:55

Question:	1	2	3	4	5	6	7	8	9	Total
Points:	12	10	14	11	10	12	10	6	15	100
Score:										

Name, Surname: \_\_\_\_\_ Student ID: \_\_\_\_\_

**In each question, show all your work to receive full credit.**

1. (a) (5 points) In a group of 20 people all of whose ages are different, in how many ways can you choose a committee of a president and two vice presidents, if the president is to be older than the vice presidents?

**Solution:** There are  $C(20,3)$  ways to choose the three persons committee. Since the oldest person is automatically the president, and the remaining two are the vice presidents, the duties are immediately assigned.

- (b) (7 points) Consider a group of 12 males and 8 females to construct a similar committee as above. This time, you have to make sure that one vice president is selected from each gender. Moreover, the president doesn't have to be older than the vice presidents anymore. In how many ways can you build the committee?

**Solution:** Now there are two ways to establish such a committee: 2 males and 1 female, or 1 male and 2 females. We know that the total number of choosing 3 persons out of 20 is  $C(20,3)$ . If we subtract from this the number of committees with no males and no females, then we will obtain a sum of the cases above. In other words,

$$MMF + MFF = C(20,3) - C(12,3) - C(8,3) = 1140 - 220 - 56 = 864$$

Clearly, within each triplet, there will be one person from one gender, and two persons from the other gender. Hence, the president must be selected among

the gender with two members, giving us 2 options in each triplet. Therefore the answer is  $864 * 2 = 1728$ .

2. (10 points) How many terms are there in the expansion of  $(a + b + c)^{10}$ ?

**Solution:** Let's assume that a term of the expansion is  $a^k b^l c^m$ . Then,  $k+l+m = 10$ . The question is, in how many different ways can we assign positive integers to  $k$ ,  $l$  and  $m$ ? We can think of 10 beads which needs to be separated by 2 sticks. The first group becomes  $k$ , the second group becomes  $l$  and the third group becomes  $m$ . For example:

0 0 0 I 0 0 0 0 0 I 0 0

means,  $k = 3$ ,  $l = 5$ ,  $m = 2$ . There are 12 places where we can place the sticks (Is). For two sticks this gives us  $C(12,2)$  possible placings. That is  $12*11/2 = 66$ . Therefore, the expansion has 66 terms.

Alternative solution: try to calculate the number of terms in the expansion of  $(a + b + c)^i$  from  $i=0$

$i=0 : 1$ ,  $i=1 : 3$ ,  $i=2 : 6$ ,  $i=3 : 10$ ,  $i=4 : 15...$

You see the pattern (add 2, add 3, add 4, add 5...). Now find for  $i=10$  by adding the numbers:  $1+2+3+4+5+6+7+8+9+10+11 = 66$ .

3. In all the questions below, balls are drawn without replacement.
- (a) (4 points) A jar contains 6 red balls, 4 blue balls, and 5 green balls. If two balls are randomly drawn, what is the probability that both balls are green?
- (b) (5 points) For the same jar as above, if three balls are drawn randomly, what is the probability that we get one ball of each color?
- (c) (5 points) Consider having two jars: one with 6 red, 4 blue, 5 green balls, the other with 4 red, 5 blue, 6 green balls. If you draw one ball from each, what is the probability that you will draw the same color from both jars.

**Solution:** a)  $\frac{5}{15} \times \frac{4}{14} = \frac{2}{21}$

b) To find the probability of drawing one ball of each color when three balls are drawn randomly, let's break it down:

Total number of balls in the jar =  $6+4+5=15$

The number of ways to choose one ball of each color:

Choose one red ball out of 6 red balls:  $C(6,1)=6$  ways

Choose one blue ball out of 4 blue balls:  $C(4,1)=4$  ways

Choose one green ball out of 5 green balls:  $C(5,1)=5$  ways

Name:

Total number of ways to select one ball of each color =  $6 \times 4 \times 5 = 120$  ways  
 Now, let's calculate the total number of ways to draw 3 balls out of 15:  
 Total number of ways to draw 3 balls out of 15 =  $C(15, 3) = \frac{15!}{3!(15-3)!} = 455$  ways  
 Therefore, the probability of getting one ball of each color when three balls are drawn randomly is =  $\frac{120}{455} = 0.2637$ .  
 c) Red from Jar 1:  $C(6,1)/C(15,1) = 6/15 = 2/5$   
 Red from Jar 2:  $4/15$   
 Probability of red from both jars:  $2/5 * 4/15 = 8/75$ .  
 Similarly, probability of blue from both jars:  $4/15 * 5/15 = 4/45$ .  
 Finally, probability of green from both jars:  $5/15 * 6/15 = 2/15$ .  
 We sum them together to find the requested probability:  
 $8/75 + 4/45 + 2/15 = 74/225$

4. Let  $a_n$  be the number of ways to climb  $n \geq 2$  stairs if the person climbing the stairs can take two stairs or three stairs at a time.
- (4 points) Find a recurrence relation for  $a_n$ . Explain your answer.
  - (4 points) What are the initial conditions?
  - (3 points) By using the recurrence relation in part (a), calculate how many ways this person can climb a flight of 10 stairs. Show your work.

**Solution:**

- $a_n = a_{n-2} + a_{n-3}$
- $a_2 = 1, a_3 = 1, a_4 = 1$ .
- $a_5 = a_2 + a_3 = 1 + 1 = 2, a_6 = a_3 + a_4 = 1 + 1 = 2, a_7 = 1 + 2 = 3, a_8 = 2 + 2 = 4, a_9 = 2 + 3 = 5, a_{10} = 3 + 4 = 7$ .

5. (10 points) Use proof by induction to show that for any positive integer  $n$ ,  $13 \cdot 7^{n-1} + 5^{2n-1}$  is divisible by 18.

**Solution:** Let  $P_n$  be the predicate that  $13 \cdot 7^{n-1} + 5^{2n-1}$  is divisible by 18.

- Base step: We show that  $P_1$  is true.  
 Since  $13 \cdot 7^{1-1} + 5^{2 \cdot 1 - 1} = 13 + 5 = 18$ ,  $P_1$  is true.

Name:

(b) Inductive step: Show that  $P_{n-1} \rightarrow P_n$ , using the inductive hypothesis that  $P_{n-1}$  is true.

We assume that  $13 \cdot 7^{n-2} + 5^{2n-3}$  is divisible by 18. (Inductive Hypothesis)

Let us rewrite  $13 \cdot 7^{n-1} + 5^{2n-1}$  as  $13 \cdot 7^{n-1} + (18+7)5^{2n-3} = 7(13 \cdot 7^{n-2} + 5^{2n-3}) + 18 \cdot 5^{2n-3}$ . The last term has a factor of 18 and the first term is divisible by 18, by the inductive hypothesis.

6. Suppose that an experiment consists of picking at random a bit string of length 5. Consider the following events to answer the questions below:

$E_1$ : the bit string chosen *begins* with 1;

$E_2$ : the bit string chosen *ends* with 1;

$E_3$ : the bit string chosen has *exactly* three 1's.

(a) (6 points) Find  $p(E_1|E_3)$ .

(b) (6 points) Determine whether  $E_1$  and  $E_2$  are independent.

**Solution:** Let the sample space  $S$  be the set of all outcomes. Let us first calculate the cardinalities of the relevant events using product rule:  $|S| = 2^5$ ,  $|E_1| = |E_2| = 2^4$ ,  $|E_3| = C(5, 3) = (5 \cdot 4 \cdot 3)/(3 \cdot 2) = 10$ , yielding  $p(E_1) = 1/2$ ,  $p(E_2) = 1/2$ ,  $p(E_3) = 10/32 = 5/16$ .

(a) (6 points)  $E_1 \cap E_3$  are the bitstrings in  $S$  beginning with 1 and having exactly two more positions with 1's. Thus,  $|E_1 \cap E_3| = C(4, 2) = (4 \cdot 3)/2 = 6$  and  $p(E_1 \cap E_3) = 6/32 = 3/16$ .

$$p(E_1|E_3) = \frac{p(E_1 \cap E_3)}{p(E_3)} = \frac{3/16}{5/16} = \frac{3}{5}.$$

(b) (6 points) For being independent, we need to check whether  $p(E_1 \cap E_2) = p(E_1)p(E_2)$ .  $E_1 \cap E_2$  contains those bitstrings in  $S$ , beginning with 1 and ending with 1. By product rule,  $|E_1 \cap E_2| = 2^3$  yielding  $p(E_1 \cap E_2) = 2^{3-5} = 1/4$ . We compare this to  $p(E_1)p(E_2) = (1/2) \cdot (1/2)$  and see they are the same. So,  $E_1$  and  $E_2$  are independent.

7. (10 points) Let  $S$  be a set of five positive integers, the maximum of which is at most 9. Prove that the sums of the elements in all the nonempty subsets of  $S$  cannot be distinct.

**Solution:** Consider the subsets of  $A$  of  $S$  where  $1 \leq |A| \leq 3$ . Since  $|S| = 5$ ,  $\binom{5}{1} + \binom{5}{2} + \binom{5}{3} = 25$  subsets  $A$ . Assume that  $s_A$  is the sum of the elements in  $A$ . So

$1 \leq s_A \leq 7 + 8 + 9 = 24$ . Using the Pigeonhole Principle, there are two subsets of  $S$  whose elements yield the same sum.

8. (6 points) The letters in the word BLUBBERER are scrambled at random (all rearrangements having equal probability); what is the probability that the two R's are adjacent to each other in the scrambled word?

**Solution:** Assume the pair of R's as a single letter. The probability:  
 $\binom{8}{(3,2,1,1,1)} / \binom{9}{(3,2,2,1,1)} = 2/9$

9. (a) (7 points) How many integer solutions  $x_1, x_2, x_3, x_4$  are there for  $x_1 + x_2 + x_3 + x_4 = 5$  where  $-1 \leq x_i$  for each  $i$ ?

**Solution:** Assume  $y_i = x_i + 1$ , where  $y_i \in N$ . Thus equation becomes  $y_1 + y_2 + y_3 + y_4 = 9$ . The total number of integer solutions are  $\binom{12}{9} = 220$

- (b) (8 points) For which positive integer  $n$  will the equations

$$x_1 + x_2 + x_3 + \dots + x_{19} = n \tag{1}$$

$$y_1 + y_2 + y_3 + \dots + y_{64} = n \tag{2}$$

have the same number of positive integer solutions?

**Solution:** For the first equation: We need the number of nonnegative integer solutions for  $w_1 + w_2 + w_3 + \dots + w_{19} = n - 19$  where  $w_i \geq 0$  for all  $1 \leq i \leq 19$ . This makes  $\binom{19+(n-19)-1}{(n-19)} = \binom{n-1}{(n-19)}$ .

For the second equation:  $z_1 + z_2 + z_3 + \dots + z_{64} = n - 64$ . This makes  $\binom{64+(n-64)-1}{(n-64)} = \binom{n-1}{(n-64)}$

Thus,  $\binom{n-1}{(n-64)} = \binom{n-1}{(n-19)} = \binom{n-1}{(64)}$ ,  $n - 19 = 63$ ,  $n = 82$