

Name: _____

BBM 205, Spring 2015

11.3.2015

Midterm I

(2 pts)

1. The notation $\exists! x P(x)$ denotes

"There exists a unique x such that $P(x)$ is true."

What are the truth values of these statements?

a) $\exists! x P(x) \rightarrow \exists x P(x)$

b) $\forall x P(x) \rightarrow \exists! x P(x)$

(2 pts)

2. Prove that if n is a positive integer, then n is even

if and only if $7n+4$ is even.

(c)

(a)

(b)

(d)

Show that this conditional statement is a tautology

$$(p \rightarrow q) \leftrightarrow (p \wedge q)$$

(2 pts)

3. Let p, q and r be the propositions

p : You get an A on the final exam

q : You do every exercise in this book

r : You get an A in this class

Write these propositions using p, q and r and logical connectives.

a) You get an A in this class, but you do not do every exercise in this book.

b) You get an A on the final exam, you do every exercise in this book, and you get an A in this class.

c) To get an A in this class, it is necessary for you to get an A on the final exam.

d) Getting an A on the final exam and doing every exercise in this book is sufficient for getting an A in this class.

a)

c)

b)

d)

(2 pts)

4. Show that this conditional statement is a tautology by using truth table:

$$(p \wedge q) \rightarrow (p \rightarrow q)$$

(3pts)

7. State the converse, contrapositive, and inverse of each of these conditional statements.

a) When I stay up late, it is necessary that I sleep until noon.

b) A positive integer is a prime only if it has no divisors other than 1 and itself.

(3pts)

d. Determine the truth value of the statement $\forall x (x^2 = 1)$

if the domain for the variables consists of

a) the nonzero real numbers

b) the nonzero integers

c) the positive real numbers

(3pts)

8. Let $Q(x)$ be the statement " $x+1 > 2x$ ". If the domain consists of all integers, what are these truth values?

a) $Q(-1)$

b) $Q(1)$

c) $\forall x \neg Q(x)$

d) $\exists x Q(x)$

e) $\forall x Q(x)$

f) $\exists x \neg Q(x)$

(2 pts)

9. Find a counterexample, if possible, to these universally quantified statements, where the domain for all variables consists of all real numbers.

a) $\forall x (x^2 \neq x)$ b) $\forall x (|x| > 0)$ c) $\forall x (x^2 \neq 2)$