

Name:

BBM 205, Spring 2015 11.3.2015

Midterm I

SOLUTIONS

(2 pts)

1. The notation  $\exists!x P(x)$  denotes

Logical "There exists a unique  $x$  such that  $P(x)$  is true."

What are the truth values of these statements?

a)  $\exists!x P(x) \rightarrow \exists x P(x)$  True

b)  $\forall x P(x) \rightarrow \exists!x P(x)$  False

(2 pts) 2. Prove that if  $n$  is a positive integer, then  $n$  is even if and only if  $7n+4$  is even.

$n, \text{even} \rightarrow 7n+4, \text{even}$ :

Since  $n=2k$  for some  $k$ , integer

$$7n+4 = 7 \cdot 2k + 4 = 2(7k+2) \text{ also even.}$$

$7n+4 \text{ even} \rightarrow n, \text{even}$

$$7n+4 = 2m \text{ where } m, \text{integer}$$

$$7n = 2m - 4 = 2(m-2), \text{even}$$

Since  $2 \nmid 7$ ,  $2 \mid n$ .  $n$  is even.

(2 pts)

3. Let  $p$ ,  $q$  and  $r$  be the propositions

$p$ : You get an A on the final exam

$q$ : You do every exercise in this book

$r$ : You get an A in this class

Write these propositions using  $p$ ,  $q$  and  $r$  and logical connectives.

a) You get an A in this class, but you do not do every exercise in this book.

b) You get an A on the final exam, you do every exercise in this book, and you get an A in this class.

c) To get an A in this class, it is necessary for you to get an A on the final exam.

d) Getting an A on the final exam and doing every exercise in this book is sufficient for getting an A in this class.

a)  $r \wedge \bar{q}$

c)  $r \rightarrow p$

b)  $p \wedge q \wedge r$

d)  $(p \wedge q) \rightarrow r$

(2 pts)

4. Show that this conditional statement is a tautology by using truth table:

$$(p \wedge q) \rightarrow (p \rightarrow q)$$

$P$	$q$	$p \wedge q$	$p \rightarrow q$	$(p \wedge q) \rightarrow (p \rightarrow q)$
0	0	0	1	1
1	0	0	0	1
0	1	0	1	1
1	1	1	1	1

(1 pt)

5. Prove that if  $x$  is rational and  $x \neq 0$ , then

$\frac{1}{x}$  is rational.

If  $x = \frac{a}{b}$  for two integers  $a, b$  with  $\gcd(a, b) = 1$ ,

then  $\frac{1}{x} = \frac{b}{a}$  also rational.

(3 pts)

6. Determine the truth value of the statement  $\forall x \exists y (xy=1)$

if the domain for the variables consists of

- a) the nonzero real numbers
- b) the nonzero integers
- c) the positive real numbers.

a)  $\forall x, \exists y = \frac{1}{x}$  so that  $xy=1$ . True

b) If  $x=2$ ,  $y=\frac{1}{2}$  so that  $xy=1$ . But  $y \notin \mathbb{Z}^+$   
False

c)  $\forall x, y = \frac{1}{x}$  so that  $xy=1$ . True

(3pts)

7. State the converse, contrapositive, and inverse of each of these conditional statements.

a) When I stay up late, it is necessary that I sleep until noon.  $\left. \begin{array}{l} q \\ p \end{array} \right\} p \rightarrow q$

b) A positive integer is a prime only if it has no divisors other than 1 and itself.  $\left. \begin{array}{l} p \\ q \end{array} \right\} p \rightarrow q$

**Converse**  $(q \rightarrow p)$  a) If I sleep... then I stay up late.

b) If no divisors other than 1... then prime.

**Contrapositive**  $(\bar{p} \rightarrow \bar{q})$  a) If I do not sleep..., then I do not stay up late.  
b) If there is any divisor other than 1 and itself, then a positive integer is not a prime.

**Inverse** a) If I do not stay up late, then I do not sleep...

b) If a positive integer is not a prime, then it has some divisor other than 1 and itself.

(3pts)

8. Let  $Q(x)$  be the statement " $x+1 > 2x$ ". If the domain consists of all integers, what are these truth values?

a)  $Q(-1)$       b)  $Q(1)$       c)  $\forall x \exists Q(x)$

d)  $\exists x Q(x)$       e)  $\forall x Q(x)$       f)  $\exists x \forall Q(x)$

a)  $-1+1=0 > -2$       b)  $1+1 \not> 2 \cdot 1$       c) As in (a),  $\exists x Q(x)$   
True                          False                          where  $x = -1$ , so, false

d)  $Q(-1)$  is true as in (a).      e)  $Q(1)$  is false as in (b).      f)  $Q(1)$  is false as in (b).  
True                          False                          (b).      True

(2 pts)

9. Find a counterexample, if possible, to these universally quantified statements, where the domain for all variables consists of all real numbers.

a)  $\forall x (x^2 \neq x)$    b)  $\forall x (|x| > 0)$    c)  $\forall x (x^2 \neq 2)$

a) False.

If  $x = 0$ ,  $x^2 = x$

b) False.

If  $x = 0$ ,

$|x| \neq 0$

c) False.

If  $x = \sqrt{2}$ ,

$x^2 = 2$ .