

Name:

BBM 205 - Spring 2015

Exam 2

SOLUTIONS

(4 points)

1. Let $P(n)$ be the statement that $1^3 + 2^3 + \dots + n^3 = \left(\frac{n(n+1)}{2}\right)^2$

for the positive integer n .

a) What is the statement $P(1)$?

b) Show that $P(1)$ is true, completing the basis step of the proof.

c) What is the inductive hypothesis?

d) Complete the inductive step.

a) $P(1): 1^3 = \left(\frac{1 \cdot 2}{2}\right)^2 = 1$ True

b) See (a).

c) I. H. : For all $i \leq n$, assume that $P(i)$ is true.

d) By (c), we assume that $P(n)$ is true.

Show that $P(n+1)$ is true.

$$1^3 + 2^3 + \dots + n^3 + (n+1)^3 \stackrel{\substack{\text{By (c)} \\ \text{(I. H.)}}}{=} \left(\frac{n(n+1)}{2}\right)^2 + (n+1)^3 =$$

$$= (n+1)^2 \left[\frac{n^2}{4} + (n+1) \right] = (n+1)^2 \left[\frac{n^2 + 4n + 4}{4} \right] = \left[\frac{(n+1)(n+2)}{2} \right]^2$$

So, $P(n+1)$ is true.

(2 points)

2. Prove that 2 divides n^2+n whenever n is a positive integer. (using induction)

$$n, \text{ even} \rightarrow n^2+n, \text{ even}; \text{ for } n \geq 1.$$

Base step: $P(1)$: $1^2+1=2$ divisible by 2.

Ind. Hypo.: $P(i)$, true for all $i \leq n$.

Ind. Step: Show that $P(n+1)$ is true.

Since $P(n)$ is true $n^2+n=2k$ for some integer k .

$$(n+1)^2+(n+1) = n^2+2n+1+n+1 = n^2+3n+2 = 2k+2n+2 = 2(k+n+1) \text{ divisible by 2.}$$

(1 point)

3. What is the cardinality of each of these sets?

a) \emptyset b) $\{\emptyset\}$ c) $\{\emptyset, \{\emptyset\}\}$ d) $\{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}$

a) 0 b) 1 c) 2 d) 3

(2 points)

4. Determine whether each of these statements is true or false.

a) $x \in \{x\}$ b) $\{x\} \subseteq \{x\}$ c) $\{x\} \in \{x\}$

d) $\{x\} \in \{\{x\}\}$ e) $\emptyset \subseteq \{x\}$ f) $\emptyset \in \{x\}$

a) True b) True c) False

d) True e) True f) False

(2 points)

5. Use the Euclidean algorithm to find $\gcd(1529, 14039)$.

$$\begin{array}{r} \text{Step 1: } 14039 \overline{)1529} \\ \underline{-13761} \\ 278 \end{array}$$

Step 2: $\gcd(1529, 14039) = \gcd(1529, 278)$

$$\begin{array}{r} 1529 \overline{)278} \\ \underline{-1390} \\ 139 \end{array}$$

Step 3: $\gcd(1529, 278) = \gcd(278, 139)$

$$\begin{array}{r} 278 \overline{)139} \\ \underline{-278} \\ 0 \end{array}$$

Step 4:

$\gcd(1529, 14039) = 139$

Answer:

$\gcd(1529, 14039) = 139$

(3 points)

6. Solve the recurrence relation with the given initial

conditions: $a_n = 2a_{n-1} + 8a_{n-2}$, $a_0 = 4$, $a_1 = 10$

$$a_n - 2a_{n-1} - 8a_{n-2} = 0$$

Let $a_n = t^n$

$$t^n - 2t^{n-1} - 8t^{n-2} = 0$$

$$t^2 - 2t - 8 = 0$$

$$(t-4)(t+2) = 0$$

$$t_1 = 4, t_2 = -2$$

Solution: $S_n = A \cdot t_1^n + B \cdot t_2^n$

where $a_0 = S_0$

$$4 = A \cdot 4^0 + B \cdot (-2)^0 = A + B \rightarrow 16 = 4A + 4B$$

$$a_1 = S_1$$

$$10 = A \cdot 4^1 + B \cdot (-2)^1 = 4A - 2B$$

$$\left\{ \begin{array}{l} 16 = 4A + 4B \\ -10 = 4A - 2B \end{array} \right. \rightarrow \begin{array}{l} 6 = 6B \rightarrow B = 1 \\ A = 3 \end{array}$$

Solution:

$$a_n = 3 \cdot 4^n + (-2)^n$$

(3 points)

7a) Find a recurrence relation and initial conditions for c_n , the minimum number of moves in which the n -disk Tower of Hanoi puzzle can be solved.

b) Solve this recurrence relation.

a) $c_{n+1} = 2c_n + 1$

b) Use substitution (or telescope) method:

$$c_{n+1} = 2c_n + 1 = 2(2c_{n-1} + 1) + 1 = 2(2(2c_{n-2} + 1) + 1) + 1 = \dots = 2(2(2(\dots(2c_1 + 1) + 1) + 1) \dots + 1) + 1 =$$

$$= 2^n + 2^{n-1} + 2^{n-2} + \dots + 1 = 2^{n+1} - 1$$

use $c_1 = 1$

(3 points)

8. Let f_n be the n^{th} Fibonacci number. Show that

$$f_0 - f_1 + f_2 - \dots - f_{2n-1} + f_{2n} = f_{2n-1} - 1 \text{ when } n \text{ is a positive integer.}$$

Use induction:

Base step: for $n=1$ $f_0 - f_1 + f_2 = f_1 - 1$ True for $n=1$

Ind. Hypo.: Assume true for all $i \leq n$ that

$$f_0 - f_1 + f_2 - \dots - f_{2i-1} + f_{2i} = f_{2i-1} - 1$$

Ind. Step: $f_0 - f_1 + f_2 - \dots - f_{2n-1} + f_{2n} - f_{2n+1} + f_{2(n+1)} =$
 $= f_{2n-1} - 1 - f_{2n+1} + f_{2n+2} = f_{2n-1} - 1 + (f_{2n+2} - f_{2n+1}) =$
 $= f_{2n} + f_{2n-1} - 1 = f_{2n+1} - 1$ true

By Ind. Hypo.