

BBM 205
Spring 2015 Exam 3

SHOW YOUR WORK TO RECEIVE FULL CREDIT.
KEEP YOUR CELLPHONE TURNED OFF.

Name: _____ SOLUTIONS

1. (4 points) For each $f(n)$ and $g(n)$, fill in the blank $f(n) = \dots(g(n))$ and $g(n) = \dots(f(n))$ with one of the symbols O, Ω or Θ . There can be more than one correct symbol.

(a) (1 point) $f(n) = 5 \log n$, $g(n) = n$; $f(n) = O(g(n))$, $g(n) = \Omega(f(n))$

(b) (1 point) $f(n) = 2^n + n^2$, $g(n) = n^3 + 3^n$; $f(n) = O(g(n))$, $g(n) = \Omega(f(n))$

(c) (1 point) $f(n) = (x^3 + 1)/(x + 1)$, $g(n) = x^2$; $f(n) = O(g(n))$, $f(n) = \Omega(g(n))$

(d) (1 point) $f(n) = n!$, $g(n) = 2^n$

So, $f(n) = \Theta(g(n))$
 $g(n) = \Theta(f(n))$

$f(n) = \Omega(g(n))$, $g(n) = O(f(n))$

2. (3 points)

Show that, for any positive integer k , $1^k + 2^k + 3^k + \dots + n^k = \Theta(n^{k+1})$ by using the definition of Θ -notation.

Step 1: Show $\sum_{i=1}^n i^k = O(n^{k+1})$

Since $1^k + 2^k + \dots + n^k \leq 1 \cdot n \cdot n^k = n^{k+1}$, true.

Step 2: Show $\sum_{i=1}^n i^k = \Omega(n^{k+1})$

$$1^k + 2^k + \dots + n^k \geq \underbrace{\left\lfloor \frac{n}{2} \right\rfloor^k + \left(\left\lfloor \frac{n}{2} \right\rfloor + 1\right)^k + \dots + (n-1)^k + n^k}_{\left\lfloor \frac{n}{2} \right\rfloor \text{ of them}} \geq \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n}{2} \right\rfloor^k \geq$$

$$\geq \frac{n^{k+1}}{2^{k+1}} = \underbrace{\frac{1}{2^{k+1}}}_{\text{a constant, since } k \text{ is fixed.}} \cdot n^{k+1}$$

By Steps 1 and 2:
 $\sum_{i=1}^n i^k = \Theta(n^{k+1})$.

3. (3 points) Find the smallest integer n such that $f(x)$ is $O(x^n)$ for each of these functions.

(a) (1 point) $f(x) = 2x^3 + x^2 \log x$, $f(x) = O(x^3)$.

(b) (1 point) $f(x) = 3x^3 + (\log x)^4$, $f(x) = O(x^3)$.

(c) (1 point) $f(x) = (x^4 + x^2 + 1)/(x^3 + 1)$, $f(x) = O(x)$.

4. (2 points) (a) (1 point) Let d be a positive integer. Use pigeonhole principle to show that among any group of $d + 1$ integers there are two with exactly the same remainder when they are divided by d .

There are $d+1$ integers (pigeons) and d remainders (pigeonholes). Since $\left\lceil \frac{d+1}{d} \right\rceil = 2$, there will always be two integers with ^{the} same remainder.

- (b) (1 point) Let n_1, n_2, \dots, n_t be positive integers. Use pigeonhole principle to show that if $n_1 + n_2 + \dots + n_t - t + 1$ objects are placed into t boxes, then for some i ($1 \leq i \leq t$), the i th box contains at least n_i objects.

If this is not true, then

box 1 contains $\leq n_1 - 1$ objects

box 2 contains $\leq n_2 - 1$ objects

\vdots

box t contains $\leq n_t - 1$ objects,

then all boxes contain $\leq (n_1 + n_2 + \dots + n_t) - t$ objects,
contradiction.

5. (2 points)

(a) (1 point) Find the coefficient of x^5 in $(4-3x)^{21}$. when $i=16$, coefficient of x^5 is

$$(4-3x)^{21} = \sum_{i=0}^{21} 4^i (-3x)^{21-i} \binom{21}{i} \quad \left\{ \begin{array}{l} \text{when } i=16, \text{ coefficient of } x^5 \text{ is} \\ \boxed{\binom{21}{16} \cdot 4^{16} \cdot (-3)^5} \end{array} \right.$$

(b) (1 point) Find the coefficient of $x^3 y^2 z^5$ in $(x+y+z)^{10}$.

$$(x+y+z)^{10} = \sum_{i,j,k} x^i y^j z^k \binom{10}{i,j,k} \quad \left\{ \begin{array}{l} \text{When } i=3, j=2, k=5, \text{ the coeff. of } x^3 y^2 z^5 \text{ is} \\ \boxed{\binom{10}{3,2,5} = \frac{10!}{3! 2! 5!}} \end{array} \right.$$

6. (2 points) Use Pascal's identity to prove the following whenever n and r are positive integers.

Claim:

$$\sum_{k=0}^{k=r} \binom{n+k}{k} = \binom{n+r+1}{r}$$

$$\sum_{k=0}^{k=r} \binom{n+k}{k} = \binom{n}{0} + \binom{n+1}{1} + \binom{n+2}{2} + \dots + \binom{n+r}{r} = \binom{n+r+1}{r}$$

Show using Induction:

For $r=0$: $\sum_{k=0}^{k=0} \binom{n+k}{k} = \binom{n}{0} = 1$ $\binom{n+0+1}{0} = 1$ True.
(Base step)

Ind. Hypo. Assume $\sum_{k=0}^{k=i} \binom{n+k}{k} = \binom{n+i+1}{i}$ for all $i < r$.

Ind. Step: $\sum_{k=0}^{k=r} \binom{n+k}{k} = \left[\sum_{k=0}^{k=r-1} \binom{n+k}{k} \right] + \binom{n+r}{r} \stackrel{\text{By Ind. Hypo.}}{=} \binom{n+r-1+1}{r-1} + \binom{n+r}{r} \stackrel{\text{By Pascal's Identity}}{=} \binom{n+r+1}{r}$

7. (1 point) How many solutions are there to the inequality

$$x_1 + x_2 + x_3 \leq 11,$$

where x_1, x_2 , and x_3 are nonnegative integers?

The solutions for $x_1 + x_2 + x_3 \leq 11$, will also be solutions

for $x_1 + x_2 + x_3 + x_4 = 11^*$, where x_1, x_2, x_3, x_4 are nonnegative integers and $x_4 = 11 - (x_1 + x_2 + x_3)$.

The number of solutions for $*$ is $\binom{11+3}{3} = \binom{14}{3}$.

8. (1 point) Seven women and nine men are on the faculty in the computer science department at a school.

- (a) (.5 points) How many ways are there to select a committee of five members of the department if at least one woman must be on the committee?

$$\binom{16}{5} - \binom{9}{5} \quad \text{or} \quad \sum_{i=1}^5 \binom{7}{i} \binom{9}{5-i}$$

- (b) (.5 points) How many ways are there to select a committee of five members of the department if at least one woman and at least one man must be on the committee?

$$\binom{7}{1} \cdot \binom{9}{1} \cdot \binom{14}{3}$$

9. (2 points) How many ways are there to distribute five balls into seven boxes if each box must have at most one ball in it if

- (a) (.5 points) both the balls and boxes are labelled?

$$\binom{7}{5} \cdot 5! = P(7, 5) = 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \text{ ways}$$

- (b) (.5 points) the balls are labelled, but the boxes are unlabelled?

1 way

- (c) (.5 points) the balls are unlabelled, but the boxes are labelled?

$$\binom{7}{5}$$

- (d) (.5 points) both the balls and boxes are unlabelled?

1 way