

1)

In Orange County, 51% of the adults are males. (It doesn't take too much advanced mathematics to deduce that the other 49% are females.) One adult is randomly selected for a survey involving credit card usage.

- a) Find the prior probability that the selected person is a male.
- b) It is later learned that the selected survey subject was smoking a cigar. Also, 9.5% of males smoke cigars, whereas 1.7% of females smoke cigars (based on data from the Substance Abuse and Mental Health Services Administration).

Use this additional information to find the probability that the selected subject is a male.

Solution

Let's use the following notation:

M = male \bar{M} = female (or not male)
C = cigar smoker \bar{C} = not a cigar smoker.

- a. Before using the information given in part b, we know only that 51% of the adults in Orange County are males, so the probability of randomly selecting an adult and getting a male is given by $P(M) = 0.51$.
- b. Based on the additional given information, we have the following:

$P(M) = 0.51$ because 51% of the adults are males

$P(\bar{M}) = 0.49$ because 49% of the adults are females (not males)

$P(C|M) = 0.095$ because 9.5% of the males smoke cigars (That is, the probability of getting someone who smokes cigars, given that the person is a male, is 0.095.)

$P(C|\bar{M}) = 0.017$. because 1.7% of the females smoke cigars (That is, the probability of getting someone who smokes cigars, given that the person is a female, is 0.017.)

Let's now apply Bayes' theorem by using the preceding formula with M in place of A, and C in place of B. We get the following result:

$$\begin{aligned} P(M|C) &= \frac{P(M) \cdot P(C|M)}{[P(M) \cdot P(C|M)] + [P(\bar{M}) \cdot P(C|\bar{M})]} \\ &= \frac{0.51 \cdot 0.095}{[0.51 \cdot 0.095] + [0.49 \cdot 0.017]} \\ &= 0.85329341 \\ &= 0.853 \text{ (rounded)} \end{aligned}$$

2)

Proof that, if a binary tree of height h has t terminal vertices, then

$$\lg t \leq h$$

Solution

Here, log function is base 2. We want to show that $t \leq 2^h$.

Using induction, base case with $h=0$ is clear.

Assuming that the statement is true when height is $h-1$, we consider a tree with height h , call it T . Remove all terminal vertices and call this new tree T' . The height of T' is $h-1$ and by inductive hypothesis, there are at most 2^{h-1} terminal vertices. Since, the removed terminal vertices in T are the children of terminal vertices in T' , there can be at most 2^h of them.

3)

Assume that the function f , g and h take on only positive values. Prove that, if $f(n) = \Theta(g(n))$ and $g(n) = \Theta(h(n))$, then $f(n) = \Theta(h(n))$.

Solution

There are constants c_1 and c_2 such that

$$c_1 h(n) \leq f(n) \leq c_2 h(n)$$

for large enough n . The constants in both inequalities are obtained by the given assumptions. (The details are omitted here.)

4) Show that a simple graph is a tree if and only if it is connected, but the deletion of any of its edges makes the graph disconnected.

Solution

→ Direction is clear, since in any tree there is a unique path between any pair of vertices and if we remove the edge uv , u and v are disconnected.

<-- Show by contrapositive: "If not a tree, then either the graph is not connected or there is a cycle by definition." Since the graph is connected, assume there is a cycle C . Then the deletion of any edge uv on that cycle would leave the graph still connected. Because if any two vertices x and y were connected using a path P that contains the edge uv , then after deleting the edge uv , x and y are still connected by using a path in the union of P and the remaining u,v -path on C . So, the deletion of any edge does not make the graph disconnected.

5) Use mathematical induction to show that given $n+1$ non-negative integers none exceeding $2n$, there is at least one integer that divides another integer in this set.

Solution

Clearly true for $n=1$.

Assume true for all i less than or equal to n . We will prove for $n+1$ numbers not exceeding

$2(n+1)$.

Case 1: If all numbers are at most $2n$, then we are done by Ind. Hypo.

Case 2: If all numbers but one number are at most $2n$, again done by Ind. Hypo.

Case 3: If at least two numbers are greater than $2n$, then there are exactly two numbers greater than $2n$. This means $2n+1$ and $2n+2$ in this set.

If $n+1$ is also in this set, then we are done, because $n+1$ divides $2n+2$.

Otherwise, we remove $2n+2$ from the set and add $n+1$ to the set. Call this new set A . By ind. Hypo, there are two integers in A that divide each other. If one of these integers is $n+1$, then the other number divides $n+1$ and this number divides also $2n+2$, done.