

BBM 205 - Discrete Structures: Midterm 1 - ANSWERS
Date: 3.11.2015, Time: 16:00 - 17:30

Ad Soyad / Name:

Öğrenci No /Student ID:

Şube /Section:

Question:	1	2	3	4	5	6	7	8	9	Total
Points:	12	4	8	24	6	12	9	15	10	100
Score:										

1. Determine the truth value of each of these statements if the domain consists of real numbers. **Give a short explanation for each answer to receive full credit.**

(a) (3 points) $\exists x(x^3 = -1)$

Solution: True. Example, $x = -1$.

(b) (3 points) $\exists x(x^4 < x^2)$

Solution: True. Example, any $x \in (-1, 1)$.

(c) (3 points) $\forall x((-x^2) = x^2)$

Solution: False. Example, any real number x other than 0.

(d) (3 points) $\forall x(2x > x)$

Solution: False. Example, any real number x that is not positive.

2. (4 points) What is the cardinality of each of these sets?
- a) $\{a\}$
 - b) $\{\{a\}\}$
 - c) $\{a, \{a\}\}$
 - d) $\{a, \{a\}, \{a, \{a\}\}\}$

Solution: 1, 1, 2, 3

3. Determine whether each of these arguments is valid. If it is valid, show the steps of your conclusion. If it is not valid, give a logical error.
- (a) (4 points)

If n is a real number such that $n > 2$, then $n^2 > 4$.
 $n \leq 2$

$n^2 \leq 4$

Solution: This conclusion is not valid. One example is when $n = -2.01$. Although $n \leq 2$, it is not true that $n^2 \leq 4$. This particular question shows that the inverse of a true statement is not always true. Here, the inverse of the first statement is false as shown by this example.

- (b) (4 points)

If it snows today, the university will close.
 The university is not closed today.

Therefore, it did not snow today.

Solution: There are various ways to show that this conclusion is valid. Let p be that *it snows today* and q be that *the university is closed*. By modus tollens, we know that

$$\begin{array}{r} p \implies q \\ \neg q \\ \hline \neg p \end{array}$$

4. Answer the following questions. Write your answer clearly and do not calculate the any number (for example, leave 2^{10} as it is instead of calculating $2^{10} = 1024$).

- (a) (4 points) How many ways are there to travel in xyz space from the origin $(0, 0, 0)$ to the point $(12, 8, 4)$ by taking steps **two units** in the positive x , positive y , or positive z direction?

Solution: Since each step has two units, there are 6 steps in x -dir., 4 steps in y -dir. and 2 steps in z -dir. to take. Thus, the answer is the following.

$$\frac{12!}{6!4!2!}$$

- (b) (4 points) How many ways are there to pack twelve **identical** DVD's into four **distinguishable** boxes so that each box contains at least one DVD?

Solution: Initially by placing one DVD in each box, we can distribute the remaining eight DVD's into the boxes in the following number of ways.

$$\binom{8 + 4 - 1}{4 - 1} = \binom{11}{3}$$

- (c) (4 points) Find the coefficient of x^4 in $(2 + 5x)^{13}$.

Solution: By the binomial theorem, we have

$$(2 + 5x)^{13} = \sum_{i=0}^{i=13} \binom{13}{i} 2^i (5x)^{13-i}.$$

Here, the power of x is 4 only when $i = 9$. So, the answer is $\binom{13}{9} 2^9 5^4$.

- (d) (4 points) How many functions are there $f : A \rightarrow B$, where $|A| = 3$, $|B| = 7$?

Solution: Let $A = \{a_1, a_2, a_3\}$. Since we have 7 choices for each element in the string $(f(a_1), f(a_2), f(a_3))$, by product rule, the answer is 7^3 .

- (e) (4 points) How many one-to-one functions are there $f : A \rightarrow B$, where $|A| = 3$, $|B| = 7$?

Solution: Let $A = \{a_1, a_2, a_3\}$. Since the values in the string $(f(a_1), f(a_2), f(a_3))$ are pairwise different in a one-to-one function, the answer is $P(7, 3) = 7 \times 6 \times 5$.

- (f) (4 points) How many subsets with an odd number of elements does a set with 10 elements have?

Solution:

$$\sum_{i=0}^{i=4} \binom{10}{2i+1}$$

5. Determine the truth value of the statement $\exists x \forall y (x \leq y^2)$ and if the domain for x and y consists of the following sets. Give a short explanation for your answer.

- (a) (2 points) the positive real numbers

Solution: False. Let $P(x, y)$ be the statement $x \leq y^2$. The explanation requires a proof by contradiction. Assume that there is an $x = \epsilon$ that satisfies $P(x, y)$ for all y in the domain. Since $P(\epsilon, 1)$ is also true, clearly $0 < \epsilon \leq 1^2 = 1$. However, this implies that $\epsilon \not\leq (\epsilon/2)^2$ and $P(\epsilon, \epsilon/2)$ is not true, a contradiction.

- (b) (2 points) the integers

Solution: True. Let x be an integer at most 0 and $x \leq y^2$ is satisfied for all integer y .

- (c) (2 points) the nonzero real numbers

Solution: True. Let x be a negative integer and $x \leq y^2$ is satisfied for all nonzero real number y .

6. (12 points) Determine whether each of these statements is true or false.

- | | |
|------------------------------|--|
| a) $0 \in \emptyset$ | b) $\emptyset \in \{0\}$ |
| c) $\{0\} \subset \emptyset$ | d) $\emptyset \subset \{0\}$ |
| e) $\{0\} \in \{0\}$ | f) $\{\emptyset\} \subset \{\emptyset\}$ |

Solution: F, F, F, T, F, T

7. (9 points) Show that if f is a function from S to T , where S and T are finite sets with $|S| > |T|$, then there are elements s_1 and s_2 in S such that $f(s_1) = f(s_2)$.

Solution: This can be shown by using pigeonhole principle.

Let pigeons be the elements of S and pigeonholes be the values of the function f . Since there are $|S|$ pigeons and at most $|T|$ pigeonholes, the fact that $|S| > |T|$ implies that the statement is true.

8. How many bit strings of length 10 contain

- (a) (4 points) an equal number of 0's and 1's?

Solution:

$$\binom{10}{5}$$

- (b) (5 points) at most four 1's?

Solution:

$$\sum_{i=0}^{i=4} \binom{10}{i}$$

- (c) (6 points) at least four 1's, where the number of 1's is even?

Solution:

$$\sum_{i=2}^{i=5} \binom{10}{2i}$$

9. How many permutations of the letters ABCDEFGH contain

(a) (2 points) the string ED?

Solution: $7!$

(b) (2 points) the strings BA and FGH?

Solution: $5!$

(c) (2 points) the strings CAB and BED?

Solution: $4!$

(d) (2 points) the strings BCA and ABF?

Solution: 0

(e) (2 points) the strings AB, DE and GH?

Solution: $5!$
