

BBM 205 - Discrete Structures: Midterm 2
Date: 28.11.2017, Time: 16:00 - 17:30

Ad Soyad / Name:
Öğrenci No /Student ID:

Question:	1	2	3	4	5	6	7	8	9	10	Total
Points:	10	12	8	14	8	16	8	8	8	8	100
Score:											

1. (a) (6 points) Use a **combinatorial proof** to show the following equality. (Kombinatorial ispat yöntemiyle aşağıdaki eşitliği gösterin.)

$$\binom{n}{r} = \binom{n-1}{r} + \binom{n-1}{r-1}$$

Solution: Let A be a set with n elements. We can count the number of r -subsets of A in two different ways. One direct way gives that number as $\binom{n}{r}$. Indirectly, fix an element $x \in A$. Group the r -subsets into two groups X_1 and X_2 . Let X_1 contain the r -subsets containing the element x . Then, $|X_1| = \binom{n-1}{r-1}$. The remaining will be the second group X_2 that consists of r -subsets that do not contain x . Then, $|X_2| = \binom{n-1}{r}$. Thus, both the LHS and the RHS count the number of r -subsets of an n -element set, thus they are equal.

- (b) (4 points) Use the equality in the first part to show that (İlk bölümdeki eşitliği kullanarak aşağıdaki gösterin.)

$$\binom{2n}{n} = 2 \binom{2n-1}{n-1}$$

Solution: By the first part,

$$\binom{2n}{n} = \binom{2n-1}{n} + \binom{2n-1}{n-1} = \binom{2n-1}{n-1} + \binom{2n-1}{n-1}.$$

2. (a) (6 points) Find a recurrence relation for the number of ways to climb n stairs if the person climbing the stairs can take one stair or two stairs at a time. Explain your answer. (Basamak sayisi n olan bir merdiveni birer ve ikiserli adimlar atarak cikmanin kac farkli yolu oldugunu gosteren rekursif bir iliski bulun. Cevabinizi aciklayin.)

Solution: Let the number of ways to climb n stairs if the person climbing the stairs can take one stair or two stairs at a time be s_n . Then, there are two possibilities based on the start of the climbing. If the person started using climbing one stair, then this climbing to the end has s_{n-1} different ways. Similarly, if the person started using climbing two stairs, then this climbing to the end has s_{n-2} different ways. Therefore, $s_n = s_{n-1} + s_{n-2}$.

- (b) (3 points) What are the initial conditions? (Baslangic kosullari nedir?)

Solution: $s_1 = 1$ and $s_2 = 2$.

- (c) (3 points) How many ways can this person climb a flight of 11 stairs? (Basamak sayisi 11 oldugunda yukarida aciklandigi gibi basamaklari cikmanin kac farkli yolu vardir?)

Solution: By using the initial conditions, we get

$$\begin{aligned} s_3 &= 1 + 2 = 3, s_4 = 3 + 2 = 5, s_5 = 5 + 3 = 8, s_6 = 8 + 5 = 13, \\ s_7 &= 13 + 8 = 21, s_8 = 21 + 13 = 34, s_9 = 34 + 21 = 55, \\ s_{10} &= 55 + 34 = 89, s_{11} = 89 + 55 = 144. \end{aligned}$$

3. (8 points) Let s_n denote the number of n -bit strings that contain the pattern 01. Find a recurrence relation to express s_n . Justify your work. (s_n , 01'i altkatar olarak iceren n elemanli bit katarlarinin sayisi olsun. s_n 'i ifade eden rekursif bir iliski bulun.)

Solution: We group such strings of length n into two. Let group-1 contain only the strings that end with 0 and group-2 contain only the strings that end with 1. All strings in group-1 must contain the substring 01 in the first $n - 1$ positions, therefore group-1 has s_{n-1} strings. To count the strings in group-2, consider the strings that end with 1 but do not contain 01 as substring. There is only one such string, basically the string of all 1's. Thus, group-2 contains $2^{n-1} - 1$ strings. Therefore, $s_n = s_{n-1} + 2^{n-1} - 1$.

4. (a) (8 points) Use extended Euclid's algorithm to calculate the multiplicative inverse of $7 \pmod{23}$. ('Extended Euclid' algoritmasını kullanarak 7 'nin $\pmod{23}$ 'de carpıma göre tersini bulun.)

Solution: By using the algorithm, we obtain the following list of output after the iterations.

$$\begin{pmatrix} 23 & 1 & 0 \\ 7 & 0 & 1 \\ 2 & 1 & -3 \\ 1 & -3 & 10 \end{pmatrix}$$

Thus, we have $-3 \cdot 23 + 10 \cdot 7 = 1$ and the multiplicative inverse of 7 is $10 \pmod{23}$.

- (b) (6 points) Find $x \pmod{23}$ if $7x + 20 \equiv 5 \pmod{23}$. (Verilen kosulu saglayan x 'i bulun)

Solution: Since the multiplicative inverse of 7 is 10 as found in the first part, we multiply both sides by 10 and obtain

$$10 \cdot (7x + 20) \equiv 1 \cdot x + 200 \equiv x + 16 \pmod{23}.$$

Since $x + 16 \equiv 5 \pmod{23}$ is given by the question, we have $10(7x + 20) \equiv 10 \cdot 5 \pmod{23}$. This together with above gives $x + 16 \equiv 50 \pmod{23}$. When we subtract 16 from both sides, we obtain $x \equiv 50 - 16 \equiv 11 \pmod{23}$.

5. (8 points) Find the solution to the recurrence relation below with the given initial questions: (Verilen rekursif iliski ve baslangic kosullarını saglayan cozum bulun.)

$$a_n = (n + 1)a_{n-1}, \quad a_0 = 2.$$

Solution: Let us use the substitution method as below:

$$\begin{aligned} a_n &= (n + 1)a_{n-1} = (n + 1)na_{n-2} = (n + 1)n(n - 1)a_{n-3} = \dots = \\ &= (n + 1)n(n - 1) \dots 3 \cdot 2 \cdot a_0 = 2(n + 1)! \end{aligned}$$

6. (a) (4 points) Explain how the pigeonhole principle can be used to show that among 11 integers at least two must end with the same digit. (Guvercinyuvasi kuralini kullanarak her 11 tamsayi arasinda mutlaka son basamagi ayni olan iki sayi olacagini gosterin.)

Solution: Since there are 10 possible digits at the end and $11 > 10$, there will be two numbers that end with the same digit.

- (b) (4 points) Explain how the generalized pigeonhole principle can be used to show that among 91 integers, there are at least ten that end with the same digit. (Genel guvercinyuvasi kuralini kullanarak her 91 tamsayi arasinda mutlaka son basamagi ayni olan en az on sayi olacagini gosterin.)

Solution: Since there are 10 possible digits at the end and $\lceil 91/10 \rceil = 10$, there will be ten numbers that end with the same digit.

- (c) (4 points) Use generalized pigeonhole principle to find how many cards must be selected from a standard deck of 52 cards to guarantee that at least four cards of the same suit are chosen. (Genel guvercinyuvasi kuralini kullanarak 52 kartlik standart kart destesinden ayni sekilden en az dort kart cikmasi icin desteden cekilmesi gereken minimum kart sayisini bulun.)

Solution: We should find the minimum N that satisfies $\lceil N/k \rceil = 4$, where $k = 4$ is the number of possible suits. The smallest N that satisfies this equality is 13.

- (d) (4 points) Use binomial theorem to show that (Binom teoremini kullanarak gosterin.)

$$\sum_{k=0}^n 3^k \binom{n}{k} = 4^n.$$

Solution: By using the binomial theorem $(x + y)^n = \sum_{k=0}^n y^{n-k} x^k \binom{n}{k}$ for $x = 3$ and $y = 1$, we obtain this result.

7. In the following parts, please circle your final answer. (Asagida verilen maddelerde, buldugunuz cevabi yuvarlak icine alin.)

- (a) (2 points) How many different ways are there to throw 8 identical balls into 15 distinguishable bins? (Sekiz ozdes topu birbirinden farkli 15 kutuya koymanin kac yolu vardir?)

Solution:

$$\binom{8 + 15 - 1}{8}$$

- (b) (2 points) A palindrome is a string whose reversal is identical to the string. How many bit strings of length n are palindromes? (Palindrom tersinden okunusu da kendisiyle ayni olan kelimedir. Uzunlugu n olan kac palindrom vardir?)

Solution: Since the strings are palindrome, once we decide for the left half of the string, the right half is fixed and we have no choice. When n is even, there are $2^{n/2}$ different palindromes. When n is odd, there are $2^{\frac{n+1}{2}}$ different palindromes.

- (c) (2 points) How many cards must be selected from a standard deck of 52 cards to guarantee that at least three spades are selected? (En az uc maca gelmesi icin standart 52 kartlik bir desteden en az kac kart cekmek gerekir?)

Solution: In the worst case, we may pick all cards possible that are not spades. There are $13 \cdot 3 = 39$ such cards. Whatever we pick in addition to these 39 cards will be spades. So, the least number of cards to pick is 42.

- (d) (2 points) How many bit strings of length 9 over the alphabet $\{a, b, c, d\}$ have either exactly three b 's or exactly five c 's? (Karakter olarak $\{a, b, c, d\}$ 'nin kullanilabilecegi kelimelerden uzunlugu 9 olan ve tam tamina uc b'yi ya da tam tamina bes c'yi iceren kac kelime vardir?)

Solution: Let A be the set of words of length 9 that have exactly three b 's. By product rule, we have $|A| = \binom{9}{3} \cdot 3^6$, since first we pick the positions for three b 's and the remaining positions can be filled with a letter from $\{a, c, d\}$. Then, let B be the set of words of length 9 that have exactly five c 's. Similarly, we have $|B| = \binom{9}{5} \cdot 3^4$. Since the answer we need is $|A \cup B| = |A| + |B| - |A \cap B|$, we need $|A \cap B|$ as well, which is given as the set of words of length 9 that have exactly three b 's and exactly five c 's given by $|A \cap B| = \binom{9}{5} \binom{5}{3} 2^2$. Thus the answer is $|A \cup B| = \binom{9}{3} \cdot 3^6 + \binom{9}{5} \cdot 3^4 - \binom{9}{5} \binom{5}{3} 2^2$.

8. In the following parts, please circle your final answer. (Asagida verilen maddelerde, buldugunuz cevabi yuvarlak icine alin.)

- (a) (2 points) How many different words can be written by using the letters in KIZILCAHAMAM when all letters are used and the word starts and ends with A? (KIZILCAHAMAM icindeki butun harfleri kullanarak ve A harfiyle baslayip biten kac kelime yazilabilir?)

Solution: There are 12 letters in KIZILCAHAMAM and the ends of the words we pick are fixed using A's. So, we only need to count the possible choices for the 10-letter words between these two A's using the available 1 K, 2 I's, 1 Z, 1 L, 1 C, 1 A, 1 H, 2 M's. The answer is $\frac{10!}{2!2!}$.

- (b) (2 points) How many ways are there to pair up 14 students so that every student is paired up? (Herkese eslestirilmis olacak sekilde 14 ogrenci kac farkli sekilde eslestirilebilir?)

Solution:

$$\binom{14}{2} \binom{12}{2} \binom{10}{2} \binom{8}{2} \binom{6}{2} \binom{4}{2} \binom{2}{2} = \frac{14!}{2!^7}$$

- (c) (2 points) How many solutions does $x_1 + x_2 + \dots + x_k = n$ have if each x_i ($1 \leq i \leq k$) must be a positive integer (at least 1)? (Verilen esitligin icindeki her x_i bir pozitif tamsayi olacak sekilde kac farkli cozum vardir?)

Solution: Since we save one unit for each x_i , we only need to find how many different ways are there to distribute $n - k$ units into k different x_i 's. The answer is $\binom{n-k+(k-1)}{k-1} = \binom{n-1}{k-1}$.

- (d) (2 points) How many ways are there to distribute 11 distinguishable balls numbered 1 through 11 among 25 distinguishable bins? (Birbirinden farkli 11 top 25 farkli kutuya kac farkli sekilde dagitilabilir?)

Solution: For each ball, we have 25 different choices. By product rule, the answer is 25^{11} .

9. (8 points) Solve the recurrence relation

$$a_n = \sqrt{\frac{a_{n-2}}{a_{n-1}}}$$

with initial conditions $a_0 = 8$, $a_1 = 1/(2\sqrt{2})$. (Verilen baslangic kosullari kullanilarak rekursif iliski icin cozum bulun.)

Solution: By letting $b_n = \log_2 a_n$, we obtain a recurrence relation for b_n as

$$b_n = \frac{b_{n-2}}{2} - \frac{b_{n-1}}{2},$$

where $b_0 = \log_2 8 = 3$ and $b_1 = \log_2(1/(2\sqrt{2})) = -1.5$. Letting $b_n = r^n$ for some r we get $r^2 = \frac{1}{2} - \frac{r}{2}$, i.e. $2r^2 + r - 1 = 0$. The roots of this equation are $r = -1$ and $r = 0.5$. Thus, $b_n = A(-1)^n + B(0.5)^n$ for some constants A and B . By using the initial conditions, we obtain $b_0 = 3 = A + B$ and $b_1 = -1.5 = -A + B/2$. Solving for A and B gives $A = 2$ and $B = 1$. This implies that $b_n = 2(-1)^n + (0.5)^n$ and thus $a_n = 2^{b_n} = 2^{2(-1)^n + (0.5)^n}$.

10. (8 points) Solve the recurrence relation with the given initial conditions. (Verilen baslangic kosullari kullanilarak rekursif iliski icin cozum bulun.)

$$a_n = 2a_{n-1} + 8a_{n-2}, \quad a_0 = 4, \quad a_1 = 10.$$

Solution: Let $a_n = r^n$ and substitute into the recurrence relation as follows.

$$r^n = 2r^{n-1} + 8r^{n-2},$$

which gives $r^2 = 2r + 8$, i.e. $r^2 - 2r - 8 = 0$ yielding two roots $r_1 = 4$ and $r_2 = -2$. Hence, we have

$$a_n = A4^n + B(-2)^n,$$

for some constants A and B . The initial conditions provide that

$$a_0 = 4 = A + B, \quad \text{and} \quad a_1 = 10 = 4A - 2B.$$

Solving for A and B above gives $A = 3$ and $B = 1$.