

BBM 205 - Discrete Structures: Final Exam
Date: 15.1.2018, Time: 9:30 - 11:30
SOLUTION SHEET

Name:

Student ID:

Question:	1	2	3	4	5	6	7	8	9	10	Total
Points:	10	22	6	8	21	10	9	14	16	8	124
Score:											

Unless specified differently, you are expected to show all your work in your answers to receive full credit from the questions.

1. (10 points) Show by using induction on the number of vertices that every tree can be colored properly by 2 colors. (Information you can use: Every tree has at least two vertices with degree 1.)

Solution: Base step: True for trees with one and two vertices.

Inductive hypothesis (I.H.): Assume that every tree on $n - 1$ vertices can be 2-colored.

Inductive step: Assume that we removed a vertex v with degree 1 from our tree T on n vertices, call this new tree T' , which can be 2-colored by I.H.. We can add v back by connecting it to its neighbor in T and obtain a 2-coloring of T by using the 2-coloring of T' . Since v has only one neighbor in T , it uses only one color. We can color v by the unused color on $N(v)$ and obtain a 2-coloring of T .

2. In the questions below, you are not expected to explain your answer.

- (a) (3 points) How many edges are there in a graph with 8 vertices each of degree 5?

Solution: $8 \cdot 5/2 = 20$.

- (b) (4 points) How many vertices and how many edges do these graphs have: a) K_n ,
b) C_n , c) $K_{m,n}$, d) Q_n

Solution: a) $(n, n(n-1)/2)$, b) (n, n) , c) $(m+n, mn)$, d) $(2^n, n2^{n-1})$.

- (c) (3 points) For which values of n are these graphs bipartite: a) K_n , b) C_n , c) Q_n

Solution: a) only when $n = 1, 2$, b) n , even, c) for every n

- (d) (3 points) For which values of m and n does the complete graph $K_{m,n}$ have a Hamilton circuit?

Solution: When $m = n$.

- (e) (3 points) Give a list of six nonnegative integers that cannot be the degree list of a graph on six vertices.

Solution: $\{3, 3, 4, 2, 2, 3\}$

- (f) (3 points) If G is a simple graph with 15 edges and \bar{G} has 13 edges, how many vertices does G have?

Solution: We have one equation with one unknown to solve, let n be the number of vertices in such a graph G . Then, $|E(G)| + |E(\bar{G})| = 15 + 13 = 28$. Therefore, $n(n-1)/2 = 28$, which gives that $n = 8$.

- (g) (3 points) Draw all nonisomorphic simple graphs with three vertices.

Solution: They are C_3 , $K_{1,2}$, K_2 and the empty graph with no edges.

3. (6 points) Let G be a graph with n vertices and e edges. Let M be the maximum degree in G . Show that $2e/n \leq M$.

Solution: We can use proof by contradiction: assume that this is not true for some graph G with n vertices and e edges with degree list $\{d_1, d_2, \dots, d_n\}$. This means $2e/n > M$ and thus


$$(d_1 + d_2 + \dots + d_n) > nM \quad (1)$$

But we also know that for every degree $d_i \leq M$, since M is the maximum degree of G . Thus,

$$(d_1 + d_2 + \dots + d_n) \leq nM. \quad (2)$$

As seen above, the equations (1) and (2) cannot be both true, contradiction.

4. (8 points) Determine whether the following pairs of graphs are isomorphic or not. (If isomorphic, provide the isomorphism function, if not isomorphic, give an explanation.)



Solution: The pair in part (a) cannot be isomorphic, because the graph on the right has a vertex with degree 4, but the graph on the left doesn't.

The pair in part (b) are isomorphic. The isomorphism function follows as:

$$f(u_1) = v_1, f(u_2) = v_3, f(u_3) = v_5, f(u_4) = v_7, f(u_5) = v_2, f(u_6) = v_4, f(u_7) = v_6.$$

5. In the questions below, all dice and coins are fair, unless specified differently.

- (a) (3 points) What is the probability that a die comes up six when it is rolled?

Solution:

$$\frac{1}{6}$$

- (b) (3 points) What is the probability that a randomly selected day of the year is in April?

Solution:

$$\frac{30}{365}$$

- (c) (3 points) What is the probability that the sum of numbers that the sum of the numbers on two dice is even when they are rolled?

Solution:

$$\frac{1 + 3 + 5 + 5 + 3 + 1}{36} = \frac{18}{36}$$

- (d) (4 points) What is the probability that when a coin is flipped six times in a row, the number of heads is at least 3?

Solution:

$$\frac{\binom{6}{3} + \binom{6}{4} + \binom{6}{5} + \binom{6}{6}}{2^6} = \frac{20 + 15 + 6 + 1}{2^6} = \frac{42}{64}$$

- (e) (4 points) What is the probability that n precedes 1 and $n - 1$ precedes 2, when we randomly select a permutation of $\{1, 2, \dots, n\}$?

Solution: There are $4! = 16$ ways to permute these four numbers and out of these permutations, only six of them satisfy the given conditions: $(n, 1, n - 1, 2)$, $(n - 1, 2, n, 1)$, $(n, n - 1, 1, 2)$, $(n, n - 1, 2, 1)$, $(n - 1, n, 1, 2)$, $(n - 1, n, 2, 1)$. By symmetry, for each permutation, there are equal number of ways to arrange the remaining elements and choose the positions of the four numbers. Hence, the probability of this event is $6/16$.

- (f) (4 points) Suppose that E and F are events such that $Pr(E) = 0.7$ and $Pr(F) = 0.5$. Show that $Pr(E \cup F) \geq 0.7$ and $Pr(E \cap F) \geq 0.2$.

Solution: Since $Pr(E \cup F) \geq Pr(E) = 0.7$, done. On the other hand, $Pr(E \cap F) = Pr(E) + Pr(F) - Pr(E \cup F) \geq 0.7 + 0.5 - 1 = 0.2$.

6. (a) (3 points) Write Euler's formula for planar graphs.

Solution: If a graph with n vertices and e edges has a planar drawing with f faces, then $n - e + f = 2$.

- (b) (7 points) Use Euler's formula to show that $K_{3,3}$ is not planar.

Solution: Observation1: Since $K_{3,3}$ does not contain a C_3 as a subgraph, every face of its planar drawing would have at least four edges.

Observation2: Since the sum of the face lengths in a planar drawing is $2e$, by Obs. 1, $4f \leq 2e$.

By Euler's formula, we have $6 - 9 + f = 2$, hence $f = 5$. By using that in Obs.2, we get $4 \cdot 5 \leq 2 \cdot 9$, which gives $20 \leq 18$, contradiction.

7. Suppose you have balls numbered $1, \dots, n$, where n is a positive integer at least 2, inside a coffee mug. You pick a ball uniformly at random, look at the number on the ball, replace the ball back into the coffee mug, and pick another ball uniformly at random.

- (a) (2 points) What is the probability that the first ball is 1 and the second ball is 2?

Solution:

$$\frac{1}{n} \frac{1}{n} = \frac{1}{n^2}$$

- (b) (3 points) What is the probability that the second ball's number is exactly one greater than the first ball's number?

Solution: This event contains the pairs $(i, i + 1)$ for $1 \leq i \leq n - 1$. Hence, the probability is $\frac{n-1}{n^2}$.

- (c) (4 points) What is the probability that the second ball is strictly less than the first ball's number?

Solution: This event contains the pairs (i, j) , where for each j , the value of i can be a number from $\{1, \dots, j - 1\}$. Hence, the total number of pairs is $\sum_{j=1}^{j=n} (j - 1) = (n - 1)n/2$. Therefore, its probability is $\frac{(n-1)n}{2n^2} = \frac{n-1}{2n}$. One can find the cardinality of this event also by making use of the symmetry, since after excluding the pairs (i, i) , there remain equal number of pairs (i, j) for which either $i < j$ or $i > j$. Hence, this event has $(n^2 - n)/2 = (n - 1)n/2$ pairs as also calculated with a different method above.

8. In the questions below, all dice and coins are fair, unless specified differently.

- (a) (5 points) Suppose that E , F_1 , F_2 , and F_3 are events from a sample space and that F_1 , F_2 , and F_3 are mutually disjoint and their union is S . Find $Pr(F_2|E)$ if $Pr(E|F_1) = 2/7$, $Pr(E|F_2) = 3/8$, $Pr(E|F_3) = 1/2$, $Pr(F_1) = 1/6$, $Pr(F_2) = 1/2$, and $Pr(F_3) = 1/3$.

Solution: By Bayes' theorem, we have

$$Pr(F_2|E) = \frac{Pr(F_2)Pr(E|F_2)}{Pr(F_1)Pr(E|F_1) + Pr(F_2)Pr(E|F_2) + Pr(F_3)Pr(E|F_3)} = \frac{(1/2)(3/8)}{(1/6)(2/7) + (1/2)(3/8) + (1/3)(1/2)} = \frac{7}{15}.$$

- (b) (5 points) What is the expected value of the sum of the numbers that appear when a pair of fair dice is rolled. Show all your work.

Solution: Let X be the sum of the numbers on the dice. Since $E(X) = \sum_{i=2}^{i=12} i \cdot Pr(X = i)$, we have

$$E(X) = 2 \cdot \frac{1}{36} + 3 \cdot \frac{2}{36} + 4 \cdot \frac{3}{36} + 5 \cdot \frac{4}{36} + 6 \cdot \frac{5}{36} + 7 \cdot \frac{6}{36} + 8 \cdot \frac{5}{36} + 9 \cdot \frac{4}{36} + 10 \cdot \frac{3}{36} + 11 \cdot \frac{2}{36} + 12 \cdot \frac{1}{36} = 7$$

- (c) (4 points) When one red die and one black die are rolled, show that the probability of the events $A = \text{sum is } 7$ and $B = \text{red die is } 1$ are independent.

Solution: To show this, we need to show that $Pr(A \cap B) = Pr(A)Pr(B)$. Since $|A| = 6$, $|B| = 6$ and $|A \cap B| = 1$, we have $Pr(A \cap B) = 1/36$, $Pr(A) = 6/36$, and $Pr(B) = 6/36$. These give $Pr(A \cap B) = Pr(A)Pr(B)$, so A and B are independent events.

The questions in the following are to receive extra credit and you are free not to solve them.

9. Zeynep and Umut are flipping coins for fun. Zeynep flips a fair coin k times and Umut flips $n - k$ times. In total there are n coin flips.

(a) (6 points) Use a combinatorial proof to show that

$$\sum_{i=0}^{i=k} \binom{k}{k-i} \binom{n-k}{i} = \binom{n}{k}.$$

You may assume $n - k \geq k$.

- (b) (6 points) Prove that Zeynep and Umut flip the same number of heads is equal to the probability that there are a total of k heads.

10. (10 points) For a graph with n vertices, if the degree of each vertex is at least $n/2$, then G is connected.