

BBM 205 - Discrete Structures: Midterm Exam - Solutions
Date: 26.11.2018, Time: 16:30 - 18:00

Name:

Student ID:


Question:	1	2	3	4	5	6	7	8	9	Total
Points:	12	9	10	10	15	10	12	12	10	100
Score:										

1. (12 points) Use the extended Euclid algorithm to find integers x and y that satisfy

$$x \cdot 2328 + y \cdot 440 = \gcd(2328, 440).$$

Solution: $2238 = 5 \cdot 440 + 128$ $\text{rem}(2238, 440) = 128 = 2238 - 5 \cdot 440$.
 $440 = 3 \cdot 128 + 56$ $\text{rem}(440, 128) = 56 = 440 - 3 \cdot 128 = 440 - 3 \cdot (2238 - 5 \cdot 440) =$
 $16 \cdot 440 - 3 \cdot 2238$.
 $128 = 2 \cdot 56 + 16$ $\text{rem}(128, 56) = 16 = 128 - 2 \cdot 56 = (2238 - 5 \cdot 440) - 2 \cdot (16 \cdot$
 $440 - 3 \cdot 2238) = 7 \cdot 2238 - 37 \cdot 440$.
 $56 = 3 \cdot 16 + 8$ $\text{rem}(56, 16) = 8 = 56 - 3 \cdot 16 = (16 \cdot 440 - 3 \cdot 2238) - 3 \cdot (7 \cdot 2238 -$
 $37 \cdot 440) = -24 \cdot 2238 - 127 \cdot 440$.
 $16 = 2 \cdot 8 + 0$. $\text{rem}(16, 8) = 0$.
By this algorithm, $x = -24$ and $y = 127$.

2. (9 points) Let G be the graph below. Carefully explain why $\chi(G) = 4$.



Solution: We use proof by contradiction, assume that we can color G with less than four colors, say using three colors as $\{1, 2, 3\}$. Let f be the color function for the vertex set as $f : V(G) \rightarrow \{1, 2, 3\}$.

Since c and d are neighbors, they must have different colors, say i and j , $i, j \in \{1, 2, 3\}$. Then, $f(x) \neq i, j$ and $f(z) \neq i, j$, because they are both neighbors of c and d . Since, there is only one color left in $\{1, 2, 3\}$ other than i and j , call this k , both $f(x) = k$ and $f(z) = k$. So, $f(x) = f(z)$.

By the similar idea (because of symmetry), we also see that $f(x) = f(y)$. However, this gives $f(x) = f(z) = f(y)$ and this is not proper coloring, because y and z are neighbors. Hence, there is no proper 3-coloring of G and we need at least 4 colors.

To see 4 colors are enough, just provide any proper coloring function $f : V(G) \rightarrow \{1, 2, 3, 4\}$. One example is as $f(x) = 1$, $f(a) = f(c) = 2$, $f(b) = f(d) = 3$, $f(y) = 1$ and $f(z) = 4$.

3. (10 points) Prove that for every positive integer n ,

$$1 \cdot 2 + 2 \cdot 3 + \cdots + n(n+1) = n(n+1)(n+2)/3.$$

Solution: $P(n)$ is the statement above.

Base case: $P(1)$ is true, since $1 \cdot 2 = 1 \cdot 2 \cdot 3/3 = 2$.

Inductive step: We can assume that $P(n)$ is true (called inductive hypothesis, I.H.) to show that $P(n+1)$ is true.

$1 \cdot 2 + 2 \cdot 3 + \cdots + (n+1)(n+2)$ can be separated into two parts:

$1 \cdot 2 + 2 \cdot 3 + \cdots + n(n+1) = n(n+1)(n+2)/3$ by I.H.

Now, adding the last term $(n+1)(n+2)$ to this gives $n(n+1)(n+2)/3 + (n+1)(n+2)$ and after some arithmetic, we see that this equals $(n+1)(n+2)(n+3)/3$, we are done.

4. (10 points) Prove that 3 divides $n^3 + 2n$ whenever n is a positive integer.

Solution: $P(n)$ is the statement that 3 divides $n^3 + 2n$.

Base case: $P(1)$ is true, since 3 divides $1^3 + 2$.

Inductive step: We can assume that $P(n)$ is true (called inductive hypothesis, I.H.) to show that $P(n+1)$ is true.

By expanding the power, we get $(n+1)^3 + 2(n+1) = n^3 + 3n + 3n^2 + 1 + 2n + 2$, which is divisible by 3 if and only if $n^3 + 2n$ divisible by 3.

I.H. tells us that this is true, hence we are done.

5. (15 points) Indicate **true or false** for the following statements about the greatest common divisor, and **provide counterexamples for those that are false**.

(a) If $\gcd(a, b) \neq 1$ and $\gcd(b, c) \neq 1$, then $\gcd(a, c) \neq 1$.

Solution: False

Let $b = 15$, $a = 3$ and $c = 5$.

Although, $\gcd(3, 15) = 3$ and $\gcd(15, 5) = 5$, we see that $\gcd(3, 5) = 1$.

(b) If $a|bc$ and $\gcd(a, b) = 1$, then $a|c$.

Solution: True

(c) $\gcd(a^n, b^n) = (\gcd(a, b))^n$.

Solution: True

(d) $\gcd(1 + a, 1 + b) = 1 + \gcd(a, b)$.

Solution: False

Let $a = 3$ and $b = 9$.

We see that $\gcd(1 + 3, 1 + 9) = \gcd(4, 10) = 2$, but $1 + \gcd(3, 9) = 1 + 3 = 4 \neq 2$.

(e) If an integer linear combination of a and b equals 1, then so does some integer linear combination of a and b^2 .


Solution: True

6. (10 points) Prove that if $m + n$ and $n + p$ are even integers, where m , n and p are integers, then $m + p$ is even.

Solution: Direct proof:

Let $m + n = 2k$ and $n + p = 2m$, where k and m are integers. Then, $m + p = (m + n) + (n + p) - 2n = 2k + 2m - 2n = 2(k + m - n)$, done.

7. (12 points) Determine if the two graphs below are isomorphic. If they are, provide an isomorphism function. If not, explain why.



Solution:

- (a) Yes. The isomorphism function is as follows:

$$f(u_1) = v_2, f(u_2) = v_1, f(u_3) = v_3, f(u_4) = v_4, \\ f(u_5) = v_8, f(u_6) = v_7, f(u_7) = v_5, f(u_8) = v_6.$$

- (b) Yes. The isomorphism function is as follows (note that there can be more than one isomorphism function in this example):

$$f(u_1) = v_1, f(u_2) = v_2, f(u_3) = v_7, f(u_4) = v_8, \\ f(u_5) = v_5, f(u_6) = v_4, f(u_7) = v_3, f(u_8) = v_6.$$

8. (a) (6 points) Draw the following graphs: K_5 , Q_2 , Q_3 . How many edges and vertices does each graph have?

Solution:

- (b) (3 points) Is there a graph (not necessarily simple) with five vertices with degrees $\{3, 2, 1, 3, 2\}$. If yes, give an example. If no, explain why.

Solution: There is no such graph, because the sum of these numbers is 11, but the degree sum of a graph should always be even.

- (c) (3 points) Represent these graphs with an adjacency matrix: C_4 , P_4 .

Solution: $\begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$ and $\begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$

9. (10 points) Use proof by contradiction to show that at least 10 of any 64 days chosen must fall on the same day of the week.

Solution: Assume that we have at most 9 from each day of the week. Then the total number of days is at most $9 \cdot 7 = 63$, but there are 64 days, a contradiction.