

BBM 205 - Discrete Structures: Midterm 2 Solutions

Date: 10.12.2019, Time: 16:00 - 17:30

Name:

Student ID:

Question:	1	2	3	4	5	6	7	8	9	Total
Points:	6	12	12	18	8	8	7	14	15	100
Score:										

1. (6 points) According to the RSA encryption system below, decryption works if for any message m , the following is true:

$$m^{de} \equiv m \pmod{pq}$$

Use Euler's Theorem to prove that $m^{de} \equiv m \pmod{pq}$ for all messages m relatively prime to pq . (Euler's Theorem says that if k is relatively prime to n then $k^{\Phi(n)} \equiv 1 \pmod{n}$.)

RSA Public Key Encryption

Beforehand The receiver creates a public key and a secret key as follows.

1. Generate two distinct primes, p and q .
2. Let $n = pq$.
3. Select an integer e such that $\gcd(e, (p-1)(q-1)) = 1$.
The *public key* is the pair (e, n) . This should be distributed widely.
4. Compute d such that $de \equiv 1 \pmod{(p-1)(q-1)}$.
The *secret key* is the pair (d, n) . This should be kept hidden!

Encoding The sender encrypts message m to produce m' using the public key:

$$m' = m^e \pmod{n}.$$


Decoding The receiver decrypts message m' back to message m using the secret key:

$$m = (m')^d \pmod{n}.$$

Solution: By the RSA algorithm and the definition of the Euler Totient function Φ , we have $de = 1 + r(p-1)(q-1) = 1 + r\Phi(pq)$ for some r . This gives

$$m^{de} \equiv m^{1+r\Phi(pq)} \pmod{pq} \equiv m \cdot (m^{\Phi(pq)})^r \pmod{pq} \equiv m \cdot 1^r \pmod{pq}.$$

2. (12 points) For each pair of graphs below, determine whether the pair is isomorphic or not. If yes, provide an isomorphism function. Otherwise, explain why they are not isomorphic.



Solution: The first pair is isomorphic. One (from many) isomorphism function is $f(u_1) = v_1, f(u_2) = v_3, f(u_3) = v_5, f(u_4) = v_7, f(u_5) = v_2, f(u_6) = v_4, f(u_7) = v_6$. The second pair is not isomorphic, since there is a vertex with degree four in the second graph, but it is missing in the first one.

3. (12 points) Use extended Euclidean algorithm to find $\gcd(9888, 6060)$ and to write $\gcd(9888, 6060)$ as a linear combination of 9888 and 6060.


Solution:

x	y	$rem(x, y)$	$= x - q \cdot y$
9888	6060	3828	$= 9888 - 6060$
6060	3828	2232	$= 6060 - 3828 = 6060 - (9888 - 6060) = -9888 + 2 \cdot 6060$
3828	2232	1596	$= 3828 - 2232 = (9888 - 6060) - (-9888 + 2 \cdot 6060)$ $= 2 \cdot 9888 - 3 \cdot 6060$
2232	1596	636	$= 2232 - 1596 = (-9888 + 2 \cdot 6060) - (2 \cdot 9888 - 3 \cdot 6060)$ $= -3 \cdot 9888 + 5 \cdot 6060$
1596	636	324	$= 1596 - 2 \cdot 636 = (2 \cdot 9888 - 3 \cdot 6060)$ $- 2 \cdot (-3 \cdot 9888 + 5 \cdot 6060) = 8 \cdot 9888 - 13 \cdot 6060$
636	324	312	$= (-3 \cdot 9888 + 5 \cdot 6060) - (8 \cdot 9888 - 13 \cdot 6060)$ $= -11 \cdot 9888 + 18 \cdot 6060$
324	312	12	$= 324 - 312 = (8 \cdot 9888 - 13 \cdot 6060) -$ $(-11 \cdot 9888 + 18 \cdot 6060) = \mathbf{19 \cdot 9888 - 31 \cdot 6060}$
312	12	0	

We observe from the results that $\gcd(9888, 6060) = 12$ that can be written as a linear combination as $12 = 19 \cdot 9888 - 31 \cdot 6060$.

4. The distance between two vertices in a graph is the length of the shortest path between them. The diameter of a graph is the distance between two vertices that are farthest apart.

(a) (6 points) What is the diameter of the following graph? Briefly explain your answer.



Solution: By checking the distances between all vertex pairs by hand, we observe that the distance between A and G is the highest. One shortest path between A and G goes through A, B, C, E, F, G , consecutively. Hence, the diameter is 5.

(b) (6 points) What is the chromatic number of this graph? Prove it.

Solution: Because, there is a triangle (also known as C_3 and K_3), at least three colors are needed to color properly, because each vertex in a triangle has to receive different colors. And three colors are enough, hence $\chi(G) = 3$. There is a proper coloring using only three colors such as: using color 1 on A, C, F , color 2 on B, D, G, H , and color 3 on E .

(c) (6 points) Suppose that every vertex in a graph is within distance t of a vertex v . Prove that the diameter of the graph is at most $2t$.

Solution: For any two vertices x and y , we consider a shortest path between v and x , call it P_1 and a shortest path between v and y , call it P_2 . Let z be the vertex, at which P_1 and P_2 intersect and it is the closest intersection to x and y . The union of the part of P_1 between x and z and the part of P_2 between y and z give us a path between x and y . This path has length at most $t + t = 2t$. Hence any shortest x, y -path has length at most $2t$. Since, x and y could be any vertices, all pairwise distances are at most $2t$, therefore the diameter as well.

5. (8 points) Show that every planar graph has a vertex of degree at most 5.

Solution: For every planar graph G with n vertices and e edges, we know that $e \leq 3n - 6$. By this and the degree-sum formula, we get

$$2e = \sum_{v \in V(G)} \deg(v) \leq 2(3n - 6) = 6n - 12.$$

As seen here, the sum of all degrees is less than $6n$ implying that the average degree is less than 6. Therefore, there must be a vertex with degree smaller than 6 (at most 5, since degrees are integers).

6. (8 points) Prove that a graph is bipartite if and only if it is 2-colorable.

Solution: There are two parts to prove:

Assume that a graph G is bipartite with parts A and B . Then, we can obtain a proper 2-coloring, by coloring A with color 1 and B with color 2. Since every edge has one endpoint in A and another in B , no edge is monochromatic. So, G is 2-colorable. Conversely, assume that a graph H has a proper 2-coloring with colors 1 and 2. Then, every edge must have one endpoint colored with 1 and another with color 2. So, we obtain two parts such that color class 1 is one part and color class 2 is another. No edge is inside one part, since that would mean the edge is monochromatic.

7. (7 points) Let $S^k = 1^k + 2^k + \dots + (p-1)^k$, where p is an odd prime and k is a positive multiple of $p-1$. Use Fermat's Theorem to prove that $S^k \equiv -1 \pmod{p}$. (Fermat's Theorem says that $a^{p-1} \equiv 1 \pmod{p}$ if p is a prime that does not divide a .)

Solution: Since k is a positive multiple of $p-1$, we can write $k = r(p-1)$ for some positive integer r . Knowing by Fermat's Theorem that $i^{p-1} \equiv 1 \pmod{p}$ for $i = 1, 2, \dots, p-1$, we have

$$S^k \equiv 1^k + 2^k + \dots + (p-1)^k \pmod{p} \equiv (p-1)1^r \pmod{p} \equiv -1 \pmod{p}.$$

8. (a) (7 points) Prove that for any number n , if the number formed by the last two digits of n is divisible by 4 then n is divisible by 4.

Solution: Assume that n has k digits and let $n = \sum_{i=0}^{k-1} 10^i n_i$. By writing this as $n = (\sum_{i=0}^{k-1} 100 \cdot 10^i n_{i+2}) + 10n_1 + n_0$, we observe that since every term other than the last two has 100 as a factor, they are divisible by 4. And $10n_1 + n_0$ is divisible by 4, because of the property of n given in the question. Thus, n is divisible by 4.

- (b) (7 points) Prove that for any number n , if the sum of the digits of n are divisible by 3, then n is divisible by 3.

Solution: The value of n in mod 3 is $\sum_{i=0}^{k-1} 10^i n_i \pmod{3} \equiv n = \sum_{i=0}^{k-1} n_i$, because $10^i \equiv 1^i \pmod{3}$. Hence, the value of n in mod 3 is the same as the value of the sum of its digits in mod 3.

9. A tree is a connected graph with no cycles. A *leaf* in a tree is a vertex with degree 1.

- (a) (8 points) Consider a tree with $n \geq 3$ vertices. What is the largest possible number of leaves the tree could have? Prove that this maximum m is possible to achieve, and further that there cannot exist a tree with more than m leaves.

Solution: It can be at most $n-1$ realized by the graph $K_{1,n}$. A tree cannot have more than $n-1$ leaves. If we assume that every vertex is a leaf, then consider a vertex x and its neighbor y . Since they are leaves and the only neighbors of each other, this means they are not connected to any other vertex in the graph, so the graph is disconnected, contradiction.

- (b) (7 points) Prove that every tree on $n \geq 2$ vertices must have at least two leaves.

Solution: One method is to use induction on the number of vertices. Here is an alternative proof. Consider a longest path P with ends x and y . Since this path is longest possible, neither x nor y can have a neighbor outside P , because in that case we could make P longer by adding this neighbor to P , contradiction. On the other hand, neither x nor y can have more than one neighbor in P . Otherwise, the longer segment of P between x and one of its neighbors together with the additional edge between x and that neighbor creates a cycle, also a contradiction. Same holds for y . Therefore, x and y has only one neighbor and they are leaves.