

BBM 205 - Discrete Structures: Final Exam Solutions
Date: 28.1.2020, Time: 10:15 - 11:45

Name:

Student ID:

Question:	1	2	3	4	5	6	7	Total
Points:	8	25	12	10	12	15	18	100
Score:								

1. (8 points) Find the solution to the recurrence relation $a_n = 2a_{n-2}$, where the initial conditions are $a_0 = 3$ and $a_1 = 5$.

Solution: By using the telescope method, we obtain for even n ,
 $a_n = 2a_{n-2} = 2^2a_{n-4} = \dots = 2^{n/2}a_0 = 3 \cdot 2^{n/2}$.
Similarly for odd n , we get $a_n = 2a_{n-2} = \dots = 2^{(n-1)/2}a_1 = 5 \cdot 2^{(n-1)/2}$.

2. In the following questions, please give your answer without needing to evaluate the formula.

(a) (5 points) In how many ways can $3n$ students be broken up into n groups of 3?

Solution: This can be calculated by the multinomial coefficient $\binom{3n}{i_1, i_2, \dots, i_n}$, where $i_1 = i_2 = \dots = i_n = 3$ but also dividing by $n!$, since the groups are not labelled. So, the answer is

$$\frac{(3n)!}{(3!)^n n!}.$$

(b) (5 points) How many $m \times n$ matrices are there with distinct entries drawn from $\{1, \dots, p\}$, where $p \geq mn$?

Solution: By product rule, this number is $p(p-1)\dots(p-mn+1) = \frac{p!}{(p-mn)!}$

(c) (5 points) How many different paths are there in a coordinate system from $(0, 0)$ to $(5, 15)$ consisting of right-steps (which increases the first coordinate by one) and up-steps (which increases the second coordinate by one)?

Solution: Once we pick where the right-steps will be, it is determined how the path will be completed. (or instead the up-steps could be picked) $\binom{5+15}{5} = \binom{20}{5}$

(d) (5 points) How many solutions with **positive integers** are there for the inequality $x_1 + x_2 + \dots + x_8 \leq 78$.

Solution: First we save one unit for every x_i and then find the number of ways to distribute $78-8=70$ units into x_i 's. We introduce a new variable, say x_9 and count how many nonnegative solutions exist for $(x_1 - 1) + (x_2 - 1) + \dots + (x_8 - 1) + x_9 = 70$. This number is given by $\binom{70+9-1}{9-1} = \binom{78}{8}$

(e) (5 points) How many undirected simple (no loops or multiple edges) graphs are there with vertices labelled as v_1, v_2, \dots, v_n ?

Solution: For every edge, we need to decide whether it is in the graph or not. (two choices for every vertex pair). Since there are $\binom{n}{2}$ pairs, the number of graphs is $2^{\binom{n}{2}}$ by product rule.

3. We want to write words can be written by using the letters $A, C, E, G, H, K, L, M, N, T, U$. Please answer the following questions without evaluating the formula.

(a) (4 points) How many words contain the subword “MATH”?

Solution: $(11 - 4 + 1)! = 8!$

(b) (4 points) How many words contain both of the subwords “MATH” and “CLEAN”?

Solution: Since, the letter A cannot come both before T and before N, there is no such word.

(c) (4 points) How many words can be written without containing the word “MATH” and “CLEAN”?

Solution: The number of words containing “MATH” is $8!$ and the number of words containing “CLEAN” is $(11 - 5 + 1) = 7!$. Since there is no word containing both as subwords, the ones containing none of these is $11! - 8! - 7!$.

4. (10 points) Solve the recurrence relation $a_n = 7a_{n-1} - 10a_{n-2}$ with initial conditions $a_0 = 5$ and $a_1 = 16$.

Solution: Let $a_n = r^n$ and substitute into the recurrence relation as follows.

$$r^n = 7r^{n-1} - 10r^{n-2},$$

which gives $r^2 = 7r - 10$, i.e. $r^2 - 7r + 10 = 0$ yielding two roots $r_1 = 2$ and $r_2 = 5$. Hence, we have

$$a_n = A2^n + B5^n,$$

for some constants A and B . The initial conditions provide that

$$a_0 = 5 = A + B, \quad \text{and} \quad a_1 = 16 = 2A + 5B.$$

Solving for A and B above gives $A = 3$ and $B = 2$.

5. (12 points) Solve the recurrence relation $a_n = 3a_{n-1} + 2^n$ with the initial condition $a_0 = 2$.

Solution: We first solve the homogeneous version $a_n = 3a_{n-1}$, which gives $a_n = A \cdot 3^n$. Then, we obtain the particular solution as $a_n = B \cdot 2^n$. To find B , we plug this solution into the recursive relation as $B \cdot 2^n = 3B \cdot 2^{n-1} + 2^n$. Thus, $2B = 3B + 2$ implying $B = -2$. So, the particular solution is $a_n = -2 \cdot 2^n = -2^{n+1}$. Combining both the solution $a_n = A \cdot 3^n$ for the homogeneous version and the particular solution, we have $a_n = A \cdot 3^n - 2^{n+1}$. Now, we use the initial condition $a_0 = 2$ to find A : $2 = A - 2$. So, $A = 4$ and the general solution is $a_n = 4 \cdot 3^n - 2^{n+1}$.

6. Let a_n be the number of ways to climb n stairs if the person climbing the stairs can take one stair or three stairs at a time.

- (a) (5 points) Find a recurrence relation for a_n . Explain your answer.

Solution: $a_n = a_{n-1} + a_{n-3}$, since we can count the possible ways in two different groups, when we started with a one-stair step or with a three-stairs step.

- (b) (5 points) What are the initial conditions?

Solution: $a_1 = 1, a_2 = 1, a_3 = 2$.

- (c) (5 points) How many ways can this person climb a flight of 10 stairs?

Solution: $a_4 = a_1 + a_3 = 1 + 2 = 3$, $a_5 = 3 + 1 = 4$, $a_6 = 4 + 2 = 6$, $a_7 = 6 + 3 = 9$, $a_8 = 9 + 4 = 13$, $a_9 = 13 + 6 = 19$, $a_{10} = 19 + 9 = 28$.

7. Prove the following statements by using the Generalized Pigeonhole Principle. For each problem, try to identify the pigeons and the pigeonholes.

- (a) (5 points) In a room of 1000 people, there are three people who share the same birthday.

Solution: There are 366 possible days (pigeonholes) and 1000 people (pigeons). By pigeonhole principle, there are at least $\lceil 1000/366 \rceil = 3$ people having the same birthday.

- (b) (6 points) In every set of 150 integers, there exist two whose difference is a multiple of 67.

Solution: There are 150 integers (pigeons) and 67 possible values that are the remainders we get after we divide these numbers by 67 (pigeonholes). By pigeonhole principle, at least $\lceil 150/67 \rceil = 3$ numbers have the same remainder when divided by 67, so the difference of any two of these numbers is a multiple of 67.

- (c) (7 points) Jellybeans of 6 different flavors are stored in 5 jars. There are 11 jellybeans of each flavor. Prove that some jar contains at least three jellybeans of one flavor and also at least three jellybeans of some other flavor.

Solution: We use pigeonhole principle twice.

First observation:

Let pigeons be 11 jellybeans of a certain flavor X and jars be pigeonholes. Because $\lceil 11/5 \rceil = 3$, one jar contains at least three jellybeans of flavor X .

Now consider a function for each flavor X_1, X_2, \dots, X_6 and label the jars 1 through 5 such that $f(X_i) = j$ is the “jar number” ($1 \leq j \leq 5$) for flavor X_i that contains at least three jellybeans of flavor X_i . (We just showed such a jar exists for every flavor.)

Second Observation:

Let pigeons be the values $f(X_1), f(X_2), \dots, f(X_6)$ and pigeonholes be jars 1, ..., 5. Since $\lceil 6/5 \rceil = 2$, there is a jar whose number is the value of f for two different flavors, that is containing at least three jellybeans from these flavors.