

BBM205 Midterm Exam I

Time: 16:00-18:00

11 November 2020

Question:	1	2	3	4	5	6	7	Total
Points:	16	18	11	13	15	10	14	97
Score:								

1. (16 points) Assume that you have a statement $P(x)$ for a number x and you aim to show the following:

$$\exists x(P(x) \implies Q(x))$$

Which of the proof ideas would be valid to prove the above statement?

Solution:

- (a) (4 points) True:

Show there exists an x such that if $Q(x)$ is false, then $P(x)$ is false.

- (b) (4 points) False:

Assume that there exists an x such that $P(x)$ is true and $Q(x)$ is false and derive a contradiction.

- (c) (4 points) True:

Find an x such that $Q(x)$ is true or $P(x)$ is false.

- (d) (4 points) True:

Assume that for all x , $P(x)$ is true and $Q(x)$ is false, and derive a contradiction.

2. Determine by using a truth table whether the following equivalences are true or false.

- (a) (6 points)

$$(x \implies y) \equiv (x \vee \neg y)$$

- (b) (6 points)

$$(x \implies y) \equiv (\neg x \implies \neg y)$$

(c) (6 points)

$$(x \implies y) \equiv ((y \wedge x) \vee \neg x)$$

Solution: Left as an exercise.

3. (11 points) Show that the number below is irrational.

$$\sqrt{x}$$

Take the value of x depending on your meeting number as follows:

$$(\text{Meeting Number}, x) = \{(1, 11), (2, 7), (3, 5), (4, 3)\}$$

Say, if you are in meeting 1, then you will take $x=11$, and similarly for others.

Solution: The proof is exactly the same as for $x = 2$, which we covered in the lecture notes. The only difference is that instead of 2, there is x .

4. (a) (5 points) Determine the following numbers by showing all your work:

$$2^{2020} \pmod{7}, \quad 3^{2020} \pmod{7}, \quad 4^{2020} \pmod{7}$$

(b) (5 points) Determine the following numbers by showing all your work:

$$2^{2020} \pmod{8}, \quad 3^{2020} \pmod{8}, \quad 4^{2020} \pmod{8}$$

(c) (3 points) Write a necessary and sufficient condition for the multiplicative inverse of a number k to exist in modulo n .

Solution:

(a)

$$\begin{array}{llll} 2^0 \equiv 1 \pmod{7}, & 3^0 \equiv 1 \pmod{7}, & 4^0 \equiv 1 \pmod{7}, \\ 2^1 \equiv 2 \pmod{7}, & 3^1 \equiv 3 \pmod{7}, & 4^1 \equiv 4 \pmod{7}, \\ 2^2 \equiv 4 \pmod{7}, & 3^2 \equiv 2 \pmod{7}, & 4^2 \equiv 2 \pmod{7}, \\ 2^3 \equiv 1 \pmod{7}, \dots & 3^3 \equiv 6 \pmod{7}, & 4^3 \equiv 1 \pmod{7}, \\ & 3^4 \equiv 4 \pmod{7}, \\ & 3^5 \equiv 5 \pmod{7}, \\ & 3^6 \equiv 1 \pmod{7}, \dots \end{array}$$

Since the pattern $(1, 2, 4)$ repeats in $2^i \pmod{7}$ and $2020 \equiv 1 \pmod{3}$, we have $2^{2020} \pmod{7} \equiv 2^1 \pmod{7} \equiv 2$.

Since the pattern $(1, 3, 2, 6, 4, 5)$ repeats in $3^i \pmod 7$ and $2020 \equiv 4 \pmod 6$, we have $3^{2020} \pmod 7 \equiv 3^4 \pmod 7 \equiv 4$.

Since the pattern $(1, 4, 2)$ repeats in $4^i \pmod 7$ and $2020 \equiv 1 \pmod 3$, we have $4^{2020} \pmod 7 \equiv 4^1 \pmod 7 \equiv 4$.

(b)

$$\begin{aligned} 2^0 &\equiv 1 \pmod 8, & 3^0 &\equiv 1 \pmod 8, & 4^0 &\equiv 1 \pmod 8, \\ 2^1 &\equiv 2 \pmod 8, & 3^1 &\equiv 3 \pmod 8, & 4^1 &\equiv 4 \pmod 8, \\ 2^2 &\equiv 4 \pmod 8, & 3^2 &\equiv 1 \pmod 8, & 4^2 &\equiv 0 \pmod 8, \\ 2^3 &\equiv 0 \pmod 8, & 3^3 &\equiv 3 \pmod 8, & 4^3 &\equiv 0 \pmod 8, \dots \\ 2^4 &\equiv 0 \pmod 8, & & & & \dots \end{aligned}$$

So, $2^{2020} \pmod 8 \equiv 0$, and $4^{2020} \pmod 8 \equiv 0$.

Since the pattern $(1, 3)$ repeats and $2020 \equiv 0 \pmod 2$, we have $3^{2020} \pmod 8 \equiv 3^0 \pmod 8 \equiv 1$.

(c) The condition is that $\gcd(k, n) = 1$.

5. The bipartite graph G has parts A and B with labelled vertices as below:

$$A = \{10a + 1, 10a + 2, 10a + 3, 10a + 4, 10a + 5\},$$

$$B = \{10a + 6, 10a + 7, 10a + 8, 10a + 9, 10a + 10\},$$

The (undirected) edge set is defined as

$$E = \{xy : x \in A, y \in B, \gcd(x, y) = 1\}$$

In this graph, find

- (a) (5 points) a maximal matching and explain why it is maximal.
- (b) (5 points) a maximum matching.
- (c) (5 points) a matching that is maximal but not maximum (if it exists).

In this question, take the value of a depending on your meeting number as below: (Meeting Number, a) = $\{(1, 0), (2, 1), (3, 2), (4, 3)\}$ Say, if you are in meeting 2, then you will take $a=1$, and similarly for others.

Solution: Say $a = 2$. The solutions for $a = 0, 1, 3$ can be solved using the same idea.

(a) (5 points) Then the edges below comprise a maximal matching:

$$\{(21, 29), (23, 30), (25, 27)\}.$$

There are no edges between the vertices 22, 24, 26 and 28, because they have a common divisor 2. Hence, this matching is maximal.

(b) (5 points) The edges below comprise a maximum matching:

$$\{(21, 26), (22, 27), (23, 30), (24, 29), (25, 28)\}.$$

(c) (5 points) Same as in part (a).

6. (10 points) Exhibit two nonisomorphic, connected graphs with x vertices and x edges.

In this question, take the value of x depending on your meeting number as follows: (Meeting Number, x) = (1,4), (2,5), (3,6), (4,7) Say, if you are in meeting 3, then you will take $x=6$, and similarly for others.

Solution: For all parts, the cycle C_x and the cycle C_{x-1} with an edge attached to it are nonisomorphic graphs with x edges and x vertices.

7. In this question, take the value of "a" depending on your meeting number as follows:

$$(\text{Meeting Number}, a) = \{(1, 2), (2, 3), (3, 2), (4, 3)\}$$

Say, if you are in meeting 4, then you will take $a=3$, and similarly for others.

(a) (5 points) Find the smallest positive integer n , for which the following is true.

$$n! \geq a^n$$

(b) (9 points) Call the number found in the first part k . Show by using induction that

$$n! \geq a^n \text{ for all } n \geq k.$$

Solution: The solutions for $a = 2, 3$ use similar ideas, so only the solution for $a = 3$ is provided below.

(a)

$$\begin{array}{ll} 1! = 1 & 3^1 = 3 \\ 2! = 2 & 3^2 = 9 \\ 3! = 6 & 3^3 = 27 \\ 4! = 24 & 3^4 = 81 \\ 5! = 120 & 3^5 = 243 \\ 6! = 720 & 3^6 = 729 \\ 7! = 5040 & 3^7 = 2187 \end{array}$$

Hence, the smallest $n = 7$.

(b) Base step: See part (a).

Inductive Step: Use the hypothesis that $n! \geq 3^n$ to show that $(n + 1)! \geq 3^{n+1}$.

We have

$$(n + 1)! = (n + 1)n! \geq (n + 1)3^n \geq 8 \cdot 3^n \geq 3^{n+1},$$

where the first inequality holds by the inductive hypothesis $n! \geq 3^n$.