

BBM205 Final Exam

Time: 13:00-14:00

3 January 2022

Name:

Student Number:

Question:	1	2	3	4	5	6	7	8	9	Total
Points:	10	11	11	9	6	8	12	11	22	100
Score:										

1. (10 points) Determine whether the following are true or false.

- (a) f is a function from the set of all bit strings to the set of integers if $f(S)$ is the number of 0 bits in S . _____

Solution: True.

- (b) Suppose $g : A \rightarrow B$ and $f : B \rightarrow C$ where $A = B = C = \{1, 2, 3, 4\}$, $g = \{(1, 4), (2, 1), (3, 1), (4, 2)\}$, and $f = \{(1, 3), (2, 2), (3, 4), (4, 2)\}$; then $f \circ g$ is $\{(1, 1), (2, 1), (3, 2), (4, 1)\}$ _____

Solution: False.

- (c) Suppose $f : \mathbb{R} \rightarrow \mathbb{Z}$, where $f(x) = \lceil 2x - 1 \rceil$. f is one-to-one. _____

Solution: False.

- (d) Suppose $f : \mathbb{R} \rightarrow \mathbb{Z}$, where $f(x) = \lceil 2x - 1 \rceil$. f is onto. _____

Solution: True.

- (e) These sets are equal for any sets A , B , and C :
 $(A - B) \cup (A - C)$, $A - (B \cap C)$ _____

Solution: True.

- (f) These sets are equal for any sets A, B, and C:

$$A - (B \cap C), (A - B) \cap (A - C)$$

Solution: False.

- (g) The following compound proposition is satisfiable:

$$((p \rightarrow q) \wedge (q \rightarrow \neg p) \wedge (p \vee q))$$

Solution: True.

- (h) For all positive integers $n \geq 2$, K_n has a Hamilton circuit.

Solution: False.

- (i) For all positive integers $n \geq 3$, W_n has a Hamilton circuit.

Solution: True.

- (j) For all positive integers $n > 2$, K_n has an Euler circuit if n is even.

Solution: False.

2. (11 points) Find $x \pmod{73}$, if $54x + 1 \equiv 13 \pmod{73}$. Show all the computations explicitly in your solution, including the inverse of 54 modulo 73, to get full credit.

Solution: Compute inverse of 54 modulo 73 using Extended Euclidean Alg:

$$73 \ 1 \ 0$$

$$54 \ 0 \ 1$$

$$19 \ 1 \ -1$$

$$16 \ -2 \ 3$$

$$3 \ 3 \ -4$$

$$1 \ -17 \ 23$$

$$1 = -17 \cdot 73 + 23 \cdot 54$$

$$\text{Therefore, } 23 \equiv 54^{-1} \pmod{73}$$

$$54x + 1 \equiv 13 \pmod{73}$$

$$54x \equiv 12 \pmod{73}$$

$$23 \cdot 54x \equiv 23 \cdot 12 \pmod{73}$$

$$x \equiv 57 \pmod{73}$$

3. (11 points) A computer generates a code that consists of a digit, a letter, another digit and another letter, with distinct digits and letters in each code, such as 0A2B, 2B9D, 0E5B etc. The letters are selected from the set, $L = \{A, B, C, D, E, F, G, H\}$ in production. What is the minimum number of codes that must be generated in order to guarantee that **at least three** of the codes are identical?


Solution: $|Digits| = 10$
 $|L| = 8$
 N: number of possible codes.
 $N = 10 \times 8 \times 9 \times 7$ (since the digits and letters are not repeated)
 $N = 5040 * 2 + 1$ guarantees that at least 3 codes are identical.

4. (9 points) Determine whether the following graph families are bipartite or not, considering n as any positive integer. Explain your answer: a) K_n b) C_n c) Q_n

Solution: a) K_n is bipartite only for $n \leq 2$, since for other cases K_n contains a triangle (K_3), which is not bipartite. b) C_n is bipartite only for even values of n . Letting the vertices of C_n be v_1, v_2, \dots, v_n , we can label the odd-indexed vertices with color 1 and even-indexed vertices with color 2, to show bipartiteness. However, with odd n , no such 2-coloring is possible where every edge has endpoints from two different colors. c) Q_n is bipartite for all n . The bipartition can be done by coloring vertices with odd weight (number of 1's in the bitstring representation of each vertex) with color 1 and vertices with even weight with color 2.

5. (6 points) a) Draw the graph represented by the following adjacency matrix.
 b) Write the degree sum and the number of edges.

Solution:
$$\begin{bmatrix} 0 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$
 The degree sum is obtained by adding all the columns (or rows) as 14. The number of edges is half of this, as 7.



6. (8 points) Find the probability (without evaluating the formula) that a randomly generated bitstring of length 12 contains exactly 3 zeros if bits are independent and if
- a 0 bit and 1 bit are equally likely,
 - the probability that a bit is 1 is 0.3.

Solution:

a) $\binom{12}{3}0.5^{12}$ b) $\binom{12}{3}0.3^90.7^3$

7. In the following questions, no explanation is needed. Please also leave your answer as a formula without evaluating it.
- (4 points) How many ways are there to travel in three dimensional coordinate system from (0,0,0) to (6,9,12) by taking steps as one unit in the +x or +y or +z direction?

Solution: $\binom{27}{6}\binom{21}{9}\binom{12}{12}$

- (4 points) How many ways are there to place 11 distinguishable balls into 5 distinguishable bins?

Solution: 5^{11}

- (4 points) Suppose that a department has 12 women and 13 men. How many ways are there to form a committee with 7 members if it must have more women than men?

Solution: $\binom{12}{4}\binom{13}{3} + \binom{12}{5}\binom{13}{2} + \binom{12}{6}\binom{13}{1} + \binom{12}{7}$

8. (11 points) 17 players on a football team show up for a match.
- How many ways do we have to choose 11 players to take the field?

Solution: $C(17, 11)$

- (b) Consider 4 (of 17) are injured. Of the 13 players who show up, 3 of them are over 30 years old. How many ways do we have to choose 10 players to take the field if at least one of these must be older than 30?

Solution: $C(10, 9)C(3, 1) + C(10, 8)C(3, 2) + C(10, 7)C(3, 3)$

9. (22 points) Two coins are in a box. The coins look identical, but one coin is fair (1/2 of Heads), while the other coin is biased with the probability 1/4 of Heads. One of the coin is selected from the box at random, without knowing which of two it is. Call the chosen coin as "**Chosen C**"

- (a) **Chosen C** is tossed twice, showing Heads both times. Given the information, what is the probability that **Chosen C** is the fair coin?

Solution: By Bayes' Rule

$$P(\text{fair}|HH) = P(HH|\text{fair})P(\text{fair})/P(HH) = 4/5$$

- (b) Are the events "*first toss of **Chosen C** is Heads*" and "*second toss of **Chosen C** is Heads*" independent? Explain briefly. (writing just "independent" or "dependent" will not be evaluated!)

Solution: They are not independent: the first toss being Heads is the evidence in favor of the coin being the fair coin, giving information about probabilities for the second toss

- (c) Find the probability that in 10 flips of **Chosen C**, there will be exactly 3 Heads. (The coin is equally likely to be either of 2 coins; do not assume it already landed Heads twice as in (a).)

Solution: X be the number of Heads in 10 tosses.

$$\begin{aligned} P(X = 3) &= P(X = 3|\text{fair})P(\text{fair}) + P(X = 3|\text{biased})P(\text{biased}) \\ &= \binom{10}{3}(1/2)^{10}(1/2) + \binom{10}{3}(1/4)^3(3/4)^7(1/2) \\ &= \frac{1}{2}\binom{10}{3}\left(\frac{1}{2^{10}} + \frac{3^7}{4^{10}}\right) \end{aligned}$$