

# BBM205

Fall 2022-2023, Midterm Exam II

December 16, 2022, 14:00-16:00

Question:	1	2	3	4	5	6	Total
Points:	16	16	16	20	16	16	100
Score:							

Name, Surname, ID: \_\_\_\_\_ Instructor: \_\_\_\_\_

1. (16 points) Recall that the Fibonacci numbers are defined with the recursive relation  $F_n = F_{n-1} + F_{n-2}$ . Given  $F_1 = 1$  and  $F_2 = 1$ , prove with induction that  $F_n$  is even whenever  $n$  is divisible by 3. For example,  $F_3 = 2$  is even,  $F_6 = 8$  is even.

**Solution:**

- Base case:  $F_1 = 1, F_2 = 1, F_3 = 2$ , so  $F_3$  is even.
- Inductive Hypothesis: Assume  $F_{3k}$  is even.
- Inductive Step: We will show that  $F_{3(k+1)} = F_{3k+3}$  is even. There are two possible cases here:
  - $F_{3k-1}$  is even. Then,  $F_{3k+1} = F_{3k} + F_{3k-1}$  must be even, because both  $F_{3k-1}$  and  $F_{3k}$  are even. Similarly, we infer that  $F_{3k+2}$  is even, and finally  $F_{3k+3}$  is even.
  - $F_{3k-1}$  is odd. Then,  $F_{3k+1} = F_{3k} + F_{3k-1}$  must be odd, because  $F_{3k}$  is even by hypothesis and even plus odd is odd. Similarly,  $F_{3k+2} = F_{3k+1} + F_{3k}$  must be odd, since  $F_{3k}$  is even and we just showed that  $F_{3k+1}$  is odd. Finally,  $F_{3k+3} = F_{3k+2} + F_{3k+1}$  must be even, because both  $F_{3k+2}$  and  $F_{3k+1}$  are odd.

An alternative inductive step avoiding the case analysis, can be writing  $F_{3k+3}$  by substituting  $F_{3k+2} = F_{3k+1} + F_{3k}$  as  $F_{3k+3} = (F_{3k+1} + F_{3k}) + F_{3k+1} = 2F_{3k+1} + F_{3k}$ . As the first term is even by the factor 2 and the second term is even by the inductive hypothesis, this is even.

2. (16 points) Prove the following:

“If  $C$  is a subset of  $B$  but not a subset of  $A$ , then  $B$  is not a subset of  $A$ .”

**Solution:** More formally we must prove the following:

$$[(C \subseteq B) \wedge (C \not\subseteq A)] \rightarrow (B \not\subseteq A)$$

We will apply the direct proof technique to prove this. Assume that the left side of the implication is true. Therefore,  $C$  is a subset of  $B$  and  $C$  is not a subset of  $A$ . Since  $C$  is a subset of  $B$ , we can write:

$$(1) \forall x(x \in C \rightarrow x \in B)$$

Similarly, since  $C$  is not a subset of  $A$ , we can write:

$$(2) \exists x(x \in C \wedge x \notin A)$$

By existential specification of (2):

$$(3) (a \in C \wedge a \notin A)$$

By simplification,

$$(4) a \in C$$

$$(5) a \notin A$$

By universal specification of (1):

$$(6) (a \in C \rightarrow a \in B)$$

Now, (4) and (6) implies the following due to modus ponens:

$$(7) a \in B$$

Combining (5) and (7) gives us:

$$(8) a \in B \wedge a \notin A$$


And, existential generalization of (8):

$$(9) \exists x(x \in B \wedge x \notin A)$$

This is equivalent to:


$$(10) B \not\subseteq A$$

3. (a) (5 points) Draw a graph with the list of degrees 3, 1, 2, 4, 1, 3. Is your graph simple?




- (b) (5 points) Draw the graph with the given adjacency matrix.

$$\begin{bmatrix} 1 & 2 & 0 & 1 \\ 2 & 0 & 3 & 0 \\ 0 & 3 & 1 & 1 \\ 1 & 0 & 1 & 0 \end{bmatrix}$$



- (c) (6 points) Draw two non-isomorphic *simple* graphs with the set of vertex degrees as {2, 2, 2, 2, 3, 3, 3, 3}.



Name:

4. In each case, say whether or not  $R$  is an equivalence relation on the given domain and explain your answer.

- (a) (4 points)  $R = \{(x, y) \in W \times W : \text{the words } x \text{ and } y \text{ start with the same letter}\}$  where  $W$  is the set of all words in the Turkish dictionary.

**Solution:** An equivalence relation is one that is reflexive, symmetric and transitive. This relation is an equivalence relation.

- Reflexive: a word starts with the same letter as itself.
- Symmetric: If word A starts with the same letter as word B, then B starts with the same letter as A.
- Transitive: If A and B starts with the same letter and B and C also do so, then A and C starts with the same letter.

- (b) (4 points)  $R = \{(x, y) \in W \times W : \text{the words } x \text{ and } y \text{ have at least one letter in common}\}$ .

**Solution:** This relation is NOT an equivalence relation.

- Reflexive: a word shares letters with itself.
- Symmetric: If A has a common letter with B, then B has the same letter common with A.
- Not transitive: (cat, hat) and (hat, hold) does not imply (cat, hold).

- (c) (4 points)  $R = \{(x, y) \in W \times W : \text{the word } x \text{ comes before the word } y \text{ in alphabetic order}\}$ .

**Solution:** This relation is NOT an equivalence relation.

- Not reflexive: a word doesn't occur in the dictionary before itself.
- Not symmetric: if A is before B, then B is NOT before A in the dictionary.
- Transitive: If A is before B, and B is before C, then A is before C in the dictionary.

- (d) (4 points)  $R = \{(x, y) \in \mathbb{R} \times \mathbb{R} : |x| \leq |y|\}$ . ( $\mathbb{R}$  is the set of real numbers.)

**Solution:** This relation is NOT an equivalence relation.

- Reflexive: the absolute value of a real number is less than or equal to itself.

- Not symmetric: if absolute A is less than absolute B, then absolute B cannot be less than absolute A.
- Transitive: If absolute A is less than or equal to absolute B, and same holds between B and C, then absolute A is less than or equal to absolute C.

(e) (4 points)  $R = \{(x, y) \in B \times B : \text{where } B \text{ is the set of all bit strings and } x \text{ and } y \text{ have the same number of 1's.}\}$ .

**Solution:** This relation is an equivalence relation.

- Reflexive: same strings have same number of 1s.
- Symmetric: if A and B have same number of 1s, then B and A also have same number of 1s.
- Transitive: If A and B have same number of ones and B and C have same number of ones, then A and C also have same number of ones.

Name:

5. (16 points) Using strong induction, prove that the solution to the recurrence relation  $a_n = 5a_{n-1} - 6a_{n-2}$ , for all  $n \geq 2$ , is  $a_n = -2 \cdot 3^n + 3 \cdot 2^n$  for the initial conditions  $a_0 = 1$  and  $a_1 = 0$ . (Provide a formal proof to get full points).

**Solution:** Let us define  $P(n)$  as “ $a_n = -2 \cdot 3^n + 3 \cdot 2^n$  for the given recurrence relation and initial conditions.”

**Basis step:**

Let's prove that  $P(0)$  and  $P(1)$  is true.

$P(0)$  is true since  $a_0 = 1$  (given) and  $a_0 = -2 \cdot 3^0 + 3 \cdot 2^0 = 1$ .

$P(1)$  is true since  $a_1 = 0$  (given) and  $a_1 = -2 \cdot 3^1 + 3 \cdot 2^1 = 0$ .

We have proved that  $P(n)$  is true for  $n=0, 1$ .

**Inductive step:**

Let  $k$  be arbitrary and  $0 \leq j \leq k$ ; assume  $k \geq 2$ .

Our Inductive hypothesis (IH) is that “ $P(j)$  holds for all  $j, 0 \leq j \leq k$ ”,

Using IH, which states that  $[P(0) \wedge P(1) \wedge \dots \wedge P(k)]$  is true, we wish to infer  $P(k+1)$  is true.

$P(k+1)$  asserts that  $a_{k+1} = -2 \cdot 3^{k+1} + 3 \cdot 2^{k+1} = -6 \cdot 3^k + 6 \cdot 2^k$

We will try to show that we can find the same value using the given recurrence relation.

Using the recurrence relation:  $a_{k+1} = 5a_k - 6a_{k-1}$ , for  $k \geq 2$ .

Note that;

- $P(k)$  is true by IH. So,  $a_k = -2 \cdot 3^k + 3 \cdot 2^k$ .
- $P(k-1)$  is also true by IH; since  $0 \leq k-1 \leq k$ , for  $k \geq 2$ .  
So,  $a_{k-1} = -2 \cdot 3^{k-1} + 3 \cdot 2^{k-1}$ .

Hence, we can replace  $a_k$  and  $a_{k-1}$  terms with the proposed solution in the recurrence relation:

$$\begin{aligned} a_{k+1} &= 5(-2 \cdot 3^k + 3 \cdot 2^k) - 6(-2 \cdot 3^{k-1} + 3 \cdot 2^{k-1}) \\ &= -6 \cdot 3^k + 6 \cdot 2^k \end{aligned}$$

Therefore,  $P(n)$  holds for all  $n \geq 0$ .

6. Determine whether each of these functions is a bijection from  $\mathbb{R} \rightarrow \mathbb{R}$ . Explain your answer for each case.

(a) (4 points)  $f(x) = -2x + 5$

**Solution:** For a function to be a bijection, it has to be a surjection and an injection. Surjection means all items in the codomain are covered and injection means each item in the codomain is mapped by exactly one item in the domain. This is a linear function. Linear functions in  $\mathbb{R}$  are surjective and injective. Therefore, it is a bijection.

(b) (4 points)  $f(x) = 4x^2 + 5$

**Solution:** This is not a bijection, because it is not surjective. For example 4 in the codomain is not mapped by any number in the domain. It is also not injective.

(c) (4 points)  $f(x) = (x + 2)/(x + 3)$

**Solution:** This is not a bijection because it is not surjective. The value 1 in the codomain is not mapped by any value in the domain.

(d) (4 points)  $f(x) = x^3$

**Solution:** This is a bijection. All real numbers have a real cubic root (surjective) and the cubes of all real numbers are distinct (injective).